

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 December 2004 (02.12.2004)

PCT

(10) International Publication Number
WO 2004/103998 A1

(51) International Patent Classification⁷: **C07D 417/12**,
215/54, 413/12, 403/12, 407/12, 401/12, 409/12, 403/14,
407/14, A61K 31/4706, 31/4709, A61P 11/00

(21) International Application Number:
PCT/EP2004/005494

(22) International Filing Date: 19 May 2004 (19.05.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0311688.6 21 May 2003 (21.05.2003) GB
0326187.2 10 November 2003 (10.11.2003) GB

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford Middlesex UB6 0NN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BALDWIN, Ian, Robert** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **BARKER, Michael, David** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **DEAN, Anthony, William** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **ELDRED, Colin, David** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **EVANS, Brian** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **GOUGH, Sharon, Lisa** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **GUNTRIP, Stephen, Barry** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **HAMBLIN, Julie, Nicole** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **HOLMAN,**

Stuart [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **JONES, Paul** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **LINDVALL, Mika, Kristian** [FI/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **LUNNISS, Christopher, James** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **REDFERN, Tracy, Jane** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **REDGRAVE, Alison, Judith** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **ROBINSON, John, Edward** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **WOODROW, Michael** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB).


(74) Agent: **COOKE, Tracey**; GlaxoSmithKline, Corporate Intellectual Property (CN925.1), 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: QUINOLINE DERIVATIVES AS PHOSPHODIESTERASE INHIBITORS



(I)

(57) Abstract: There are provided according to the invention novel compounds of formula (I) or pharmaceutically acceptable salts thereof, wherein R1, R2, R19, R20 and R34 are as described in the specification, processes for preparing them, formulations containing them and their use in therapy for the treatment of inflammatory diseases.



Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

QUINOLINE DERIVATIVES AS PHOSPHODIESTERASE INHIBITORS


The present invention relates to quinoline compounds, processes for their preparation, intermediates usable in these processes, and pharmaceutical compositions containing the compounds. The invention also relates to the use of the quinoline compounds in therapy, for example as inhibitors of phosphodiesterases and/or for the treatment and/or prophylaxis of inflammatory and/or allergic diseases such as chronic obstructive pulmonary disease (COPD), asthma, rheumatoid arthritis or allergic rhinitis.

WO 02/20489 A2 (Bristol-Myers-Squibb Company) discloses 4-aminoquinoline derivatives wherein the 4-amino group NR^4R^5 may represent an acyclic amino group wherein R^4 and R^5 may each independently represent hydrogen, alkyl, cycloalkyl, aryl, heteroaryl etc.; NR^4R^5 may alternatively represent an aliphatic heterocyclic group. The compounds are disclosed as inhibitors of cGMP phosphodiesterase, especially type 5 (PDE5).

EP 0 480 052 (Otsuka Pharmaceutical Co. Ltd.) discloses 4-aminoquinoline-3-carboxamides wherein the 4-amino group NHR^4 may represent an amino group wherein R^4 represents phenyl, tetrahydronaphthyl or naphthyl, optionally substituted with alkyl, halogen, alkoxy etc.; and the 3-carboxamide group CONR^2R^3 represents a primary, secondary or tertiary carboxamide group. The compounds are disclosed as inhibitors of gastric acid secretion, and as cytoprotective agents; inhibition of the ATPase activated by H^+ and K^+ at the gastric wall cells is also disclosed.

It is desirable to find new compounds which bind to, and preferably inhibit, phosphodiesterase type IV (PDE4).

According to the invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein:

5 R^1 is

C_{1-6} alkyl;

10 C_{3-7} cycloalkyl or C_{3-7} cycloalkyl(C_{1-4} alkyl)- wherein the C_{3-7} cycloalkyl is optionally substituted by one or more substituents selected from =O and OH;

C_{4-7} cycloalkyl fused to an aryl ring;

15 Aryl or aryl(C_{1-6} alkyl)- wherein the aryl is optionally substituted by one or more substituents selected from C_{1-6} alkyl, C_{1-6} alkylCONR⁶-, C_{1-6} alkylCO-, halogen, -CF₃, -
(CH₂)_mOH, -OCF₃, C_{1-6} alkoxy-, C_{1-6} alkoxy(C_{1-4} alkyl)-, C_{1-6} alkoxy C_{2-6} alkoxy-, C_{1-6} alkoxycarbonyl, -CN, R^4R^5 NCO, R^7R^8 N-, R^9R^{10} NCONR¹¹-, HO(CH₂)₂₋₆O-,
 $R^{12}R^{13}$ NSO₂(CH₂)_m-, (4-morpholinyl) C_{2-6} alkoxy, -NR¹⁴SO₂ C_{1-6} alkyl, aryloxy, heteroaryl
20 (optionally substituted by C_{1-6} alkyl), CO₂H, $R^{21}R^{22}$ N(C_{1-4} alkyl)-, C_{1-6} alkoxyCONR²³(CH₂)_m-,
aryl(optionally substituted by C_{1-6} alkyl);

Aryl fused to a C_{4-7} cycloalkyl ring, wherein the cycloalkyl ring is optionally substituted by one or more =O;

25 Aryl fused to a heterocyclyl ring, wherein the heterocyclyl ring is optionally substituted by one or more substituents selected from =O, -COC₁₋₄alkyl, C_{1-4} alkyl;


Heteroaryl or heteroaryl(C₁₋₆alkyl)- wherein the heteroaryl is optionally substituted by one or more substituents selected from: C₁₋₆alkyl, aryl(C₁₋₄alkyl), C₁₋₆alkoxy, halogen, C₁₋₆alkoxyCO; or

5

Heterocyclyl optionally fused to an aryl or heteroaryl ring;

R² is hydrogen or C₁₋₆alkyl;



10 R³⁴ is hydrogen or a group of formula:



wherein R³ is

15

C₁₋₆alkyl optionally substituted by one or more substituents selected from -OH, -NR¹⁶COR¹⁵, -NR¹⁷R¹⁸, -CO₂R²⁴, C₁₋₆alkoxyCONR²⁵-, -CONR²⁶R²⁷, C₁₋₆alkoxy-, C₁₋₆alkylSO₂NR³³-, or a group having one of the following formulae;



20

C₃₋₇cycloalkyl;

25 Aryl or aryl(C₁₋₆alkyl)- wherein the aryl is optionally substituted by one or more substituents selected from C₁₋₆alkyl-, halogen-, C₁₋₆alkoxy-, -CO₂R²⁸, -CH₂CO₂H, -OH, aryl(optionally substituted by a C₁₋₆alkoxy group), heteroaryl, -CONR²⁹R³⁰, C₃₋₇cycloalkoxy, C₃₋₇cycloalkyl(C₁₋₆alkoxy)-, -CF₃;

Heteroaryl or heteroaryl(C₁₋₆alkyl)- wherein the heteroaryl is optionally substituted by one or more C₁₋₆alkyl or -CONR²⁹R³⁰ groups; or

Heterocyclyl which is optionally substituted by one of more substituents selected from


- 5 C₁₋₆alkyl-, C₁₋₆alkylCO-, C₃₋₇cycloalkylCO-, heteroarylCO- (optionally substituted by one or more C₁₋₄alkyl- groups), C₁₋₆alkoxyCO-, arylCO-, R³¹R³²NCO-, C₁₋₆alkylSO₂-, arylSO₂-, -heteroarylSO₂ (optionally substituted by one or more C₁₋₄alkyl or C₁₋₄alkylCONH- groups)
The heterocyclyl is linked to the S(=O)_n moiety through a carbon atom.

- 10 m is 0-6

n is 0, 1 or 2;

R¹⁹ is hydrogen, C₁₋₆alkyl or a group of formula:

15



R²⁰ is hydrogen, C₁₋₆alkyl, halogen or C₁₋₆alkoxy;

- 20 R⁴⁻¹⁸, R²¹⁻²⁵, R²⁸ and R³¹⁻³³ all independently represent H, C₁₋₆ alkyl;

R²⁶ and R²⁷ independently represent H, C₁₋₆ alkyl, C₃₋₇cycloalkyl or heterocyclyl;

R²⁹ and R³⁰ independently represent H, C₁₋₆alkyl optionally substituted by OH;

25

R⁷ and R⁸ together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

R⁹ and R¹⁰ together with the nitrogen atom to which they are attached may form a
30 heterocyclyl ring;

R¹⁷ and R¹⁸ together with the nitrogen atom to which they are attached may form a heterocyclyl ring such as morpholine;

5 R²¹ and R²² together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

R²⁶ and R²⁷ together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

10 R²⁹ and R³⁰ together with the nitrogen atom to which they are attached may form a heterocyclyl ring such as morpholine;

R³¹ and R³² together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

15

with the proviso that R³⁴ and R¹⁹ cannot both represent R³S(=O)_n-.

As used herein, the term "alkyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms. For example, C₁₋₆alkyl means a
20 straight or branched alkyl chain containing at least 1, and at most 6, carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, *n*-pentyl and *n*-hexyl. A C₁₋₄alkyl group is preferred, for example methyl, ethyl or isopropyl. The said alkyl groups may be optionally substituted with one or more fluorine atoms, for example, trifluoromethyl.

25

As used herein, the term "alkoxy" refers to a straight or branched chain alkoxy group, for example, methoxy, ethoxy, prop-1-oxy, prop-2-oxy, but-1-oxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy or hexyloxy. A C₁₋₄alkoxy group is preferred, for
30 example methoxy or ethoxy. The said alkoxy groups may be optionally substituted with one or more fluorine atoms, for example, trifluoromethoxy.

As used herein, the term "cycloalkyl" refers to a non-aromatic hydrocarbon ring containing the specified number of carbon atoms. For example, C₃₋₇cycloalkyl means a non-

aromatic ring containing at least three, and at most seven, ring carbon atoms. Examples of "cycloalkyl" as used herein include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. A C₃₋₆cycloalkyl group is preferred, for example cyclopentyl or cyclohexyl.

5

When used herein, the term "aryl" refers to, unless otherwise defined, a mono- or bicyclic carbocyclic aromatic ring system containing up to 10 carbon atoms in the ring system, for instance phenyl or naphthyl, optionally fused to a C₄₋₇cycloalkyl or heterocyclyl ring.

- 10 As used herein, the terms "heteroaryl ring" and "heteroaryl" refer to, unless otherwise defined, a monocyclic five- to seven- membered heterocyclic aromatic ring containing one or more heteroatoms selected from oxygen, nitrogen and sulfur. In a particular aspect such a ring contains 1-3 heteroatoms. Preferably, the heteroaryl ring has five or six ring atoms. Examples of heteroaryl rings include, but are not limited to, furyl, thienyl, pyrrolyl, 15 oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, imidazolyl, pyrazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl and triazinyl. The terms "heteroaryl ring" and "heteroaryl" also refer to fused bicyclic heterocyclic aromatic ring systems containing at least one heteroatom selected from oxygen, nitrogen and sulfur, preferably from 1-4 heteroatoms, more preferably from 1 to 3 heteroatoms. Preferably, 20 the fused rings each independently have five or six ring atoms. Examples of fused aromatic rings include, but are not limited to, quinoliny, isoquinoliny, quinazolinyl, quinoxaliny, cinnoliny, naphthyridiny, indolyl, indazolyl, pyrrolopyridiny, benzofuranyl, benzothieryl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzoxadiazolyl and benzothiadiazolyl. The heteroaryl may attach to the 25 rest of the molecule through any atom with a free valence.

- As used herein, the term "heterocyclyl" refers to a monocyclic three- to seven-membered saturated or non-aromatic, unsaturated ring containing at least one heteroatom selected from oxygen, nitrogen and sulfur. In a particular aspect such a ring contains 1 or 2 30 heteroatoms. Preferably, the heterocyclyl ring has five or six ring atoms. Examples of heterocyclyl groups include, but are not limited to, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, imidazolidinyl, pyrazolidinyl, piperazinyl, morpholiny, thiomorpholiny, diazepinyl, azepinyl, tetrahydrofuranyl, tetrahydropyranyl, and 1,4-dioxanyl.

As used herein, the terms "halogen" or "halo" refer to fluorine, chlorine, bromine and iodine. Preferred halogens are fluorine, chlorine and bromine. Particularly preferred halogens are fluorine and chlorine.

5

As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s) which occur and events that do not occur.

10 As used herein, the term "substituted" refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated. Where one or more substituents are referred to, this will refer for instance to 1 to 4 substituents, and preferably to 1 or 2 substituents.

15 In one embodiment, R¹ is selected from:

C₃₋₇ cycloalkyl, in particular cyclohexyl;

20 Aryl optionally substituted by one or more substituents selected from: C₁₋₆alkyl, halogen, C₁₋₆alkoxy, C₁₋₆alkoxy(C₁₋₄alkyl)-, -CN, -(CH₂)_mOH, -CF₃, C₁₋₆alkoxyC₂₋₆alkoxy-, R⁴R⁵NCO, C₁₋₆alkylCONR⁶-, R⁷R⁸N-, C₁₋₆alkoxycarbonyl, HO(CH₂)₂₋₆O-, C₁₋₆alkylCO-, heteroaryl (optionally substituted by C₁₋₆alkyl) particularly oxazolyl, pyrazolyl or 1,2,4-oxadiazolyl;

Aryl(C₁₋₂alkyl) wherein the aryl is optionally substituted by -OH;

25

Aryl fused to a C₅₋₆ cycloalkyl ring wherein the cycloalkyl is optionally substituted by (=O);

Aryl fused to a heterocyclyl ring, wherein the heterocyclyl ring is optionally substituted by one or more substituents selected from =O, C₁₋₄alkyl;

30


Heteroaryl optionally substituted by one or more C₁₋₆alkyl, halogen (in particular chlorine or fluorine) or C₁₋₆alkoxy groups in particular wherein heteroaryl represents benzothiazolyl, benzisoxazolyl, benzimidazolyl, indazolyl, pyridyl and pyrazolyl;

Heteroaryl(C₁₋₂alkyl) wherein the heteroaryl is optionally substituted by one or more C₁₋₆alkyl groups, in particular wherein heteroaryl represents pyridyl, pyrazolyl; or

5 Heterocyclyl, in particular tetrahydropyranyl.


Examples of suitable aryl fused to a C₅₋₆ cycloalkyl ring include:


10









Examples of suitable aryl fused to heterocyclyl rings include:

15









The following aryl fused to heterocyclyl rings are further embodiments:




5



In a further embodiment, R^1 is selected from:


- 10 Aryl optionally substituted by one or more substituents selected from: methyl, ethyl, fluorine, chlorine, -CN, -CH₂OH, -OMe, -OH, -NMe₂, -O(CH₂)₂OH, -CF₃, -COMe, 1,2,4-oxadiazolyl substituted by methyl; particular substituted aryl groups include; 3-(methyloxy)phenyl, 3-methylphenyl, 3-cyanophenyl, 3-fluorophenyl, 3-chlorophenyl, 4-fluoro-3-(methyloxy)phenyl, 3-acetylphenyl, 4-hydroxy-3-(methyloxy)phenyl, 2-fluoro-3-chlorophenyl, 2,3-difluorophenyl, 3,5-difluorophenyl;
- 15

Aryl fused to a cyclohexane or cyclopentane ring, wherein the cyclopentane ring is optionally substituted by (=O); in particular the following fused systems:



Aryl fused to a heterocyclyl ring, optionally substituted by methyl; in particular the following heterocyclyl ring fused aryl systems:

5




- 10 Heteroaryl optionally substituted by one or more methyl, ethyl, flourine, chlorine or methoxy groups; in particular a pyridyl, benzimidazolyl, pyrazolyl or indazolyl group optionally substituted by one or more methyl, ethyl, flourine, chlorine, or methoxy groups; preferably 1-methyl-1H-benzimidazolyl-6-yl, 1-methyl-1H-indazol-6-yl, 5-(methoxy)-3-pyridinyl, 3-pyridinyl, 1-ethyl-1H-pyrazol-5-yl, 5-methyl-3-pyridinyl, 1,3-
 15 benzothiazol-6-yl, 5-fluoro-3-pyridinyl, or 5-chloro-3-pyridinyl.

In one embodiment, R^2 is hydrogen.

In one embodiment R^3 is selected from:

- 20 C_{1-6} alkyl which is optionally substituted by one or more substituents selected from $-NR^{16}COR^{15}$; $OH-$, $C_{1-6}alkoxyCONR^{25}-$, $-CONR^{26}R^{27}$, $-NH_2$, $-NR^{17}R^{18}$, $-CO_2R^{24}$, $C_{1-6}alkoxy-$; or a group having one of the following formulae:



C₃₋₇cycloalkyl;

Aryl optionally substituted by one or more substituents selected from C₁₋₆alkyl-, halogen-,
 5 C₁₋₆alkoxy-, -CO₂R²⁸, -OH, -CONR²⁹R³⁰, C₃₋₇cycloalkoxy, C₃₋₇cycloalkyl(C₁₋₆alkoxy);

Aryl(C₁alkyl) wherein the aryl is optionally substituted by one or more C₁₋₆alkoxy groups;

Heteroaryl or heteroaryl(C₁₋₆alkyl) which is optionally substituted by one or more C₁₋₆alkyl
 10 or -CONR²⁹R³⁰ groups; or

Heterocyclyl which is optionally substituted by one or more substituents selected from
 C₁₋₆alkyl-, C₁₋₆alkylCO-, C₃₋₇cycloalkylCO-, heteroarylCO- (optionally substituted by one or
 more C₁₋₄alkyl- groups), C₁₋₆alkoxyCO-, arylCO-, C₁₋₆alkylSO₂-.

15

In an alternative embodiment R³ is selected from:

methyl, ethyl, n-propyl, *tert*-butyl, isopropyl, MeCONH(CH₂)₂-, Me₂NCO(CH₂)₂-;

Cyclopentyl;

20

Aryl optionally substituted by one or more methoxy, methyl, -CONH₂ or -CONMe₂
 groups; in particular: 4-(methyloxy)phenyl, phenyl, 3-[(dimethylamino)carbonyl]phenyl, 4-
 methylphenyl, 3-[(methyloxy)carbonyl]phenyl, 3,4-bis(methyloxy)phenyl, 3,4,5-
 tris(methyloxy)phenyl, 3-(ethyloxy)phenyl;

25

Heterocyclyl which is optionally substituted by one of more substituents selected from
 MeCO-, cyclopropylCO-, 2-furylCO-, or MeSO₂-; in particular wherein the heterocyclyl
 group is tetrahydropyran-4-yl; a tetrahydrofuran-3-yl; or piperidinyl substituted by one or
 more substituents selected from MeCO-, cyclopropylCO-, 2-furylCO-, or MeSO₂-,
 30 especially 1-acetyl-4-piperidinyl, 1-(2-furanylcabonyl)-4-piperidinyl, 1 -
 (cyclopropylcarbonyl)-4-piperidinyl;

Heteroaryl wherein the heteroaryl represents 3-pyridyl which is optionally substituted by CONMe₂, especially 5-[(dimethylamino)carbonyl]-3-pyridinyl.

5 In one embodiment R⁹ and R¹⁰ together with the nitrogen to which they are attached represent 4-morpholinyl.

In one embodiment R⁴, R⁵, R⁶, R⁷, R⁸, R¹¹⁻¹⁶, and R²¹⁻²⁵ and R²⁸⁻³³ are independently selected from hydrogen and methyl.

10 In one embodiment R²⁶ and R²⁷ are independently selected from hydrogen, methyl, cyclopropyl, or 4-tetrahydropyranyl; or R²⁶ and R²⁷ together with the nitrogen to which they are attached form a heterocyclyl ring, in particular pyrrolidinyl or morpholinyl.

15 In one embodiment R¹⁹ is hydrogen, C₁₋₆alkyl or MeSO₂-. In a further embodiment R¹⁹ is hydrogen or methyl, especially hydrogen.


In one embodiment R²⁰ is hydrogen, halogen or C₁₋₆alkyl. Alternatively R²⁰ is hydrogen, chlorine, fluorine, methyl or ethyl. In a further embodiment R²⁰ is methyl, ethyl or chlorine.

20 In one embodiment m is 0 or 1.

In one embodiment n is 1 or 2, especially 2.

In one embodiment R³⁴ represents a group of formula:

25



It is to be understood that the present invention covers all combinations of substituent groups referred to herein above.

30

It is to be understood that the present invention covers all combinations of particular and preferred groups described herein above.

Particular compounds according to the invention include those mentioned in the examples and their pharmaceutically acceptable salts. Specific examples which may be mentioned include:

Example 7: 4-[(3-methylphenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 8: 4-[(3-cyanophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 20: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 27: 4-[[3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 32: 4-[[4-fluoro-3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 35: 4-[(3-chlorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 43: 4-(1,3-benzothiazol-6-ylamino)-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 45: 4-[(1-methyl-1H-benzimidazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 52: 4-[(3-cyanophenyl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 66: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 74: 4-[[3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 89: 6-(cyclopentylsulfonyl)-4-[(3-fluorophenyl)amino]-3-quinolinecarboxamide,

Example 128: 4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide,

5

Example 129: 6-[(1,1-dimethylethyl)sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

Example 130: 6-[[2-(acetylamino)ethyl]sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide.

10

Example 133: 6-[(1,1-dimethylethyl)thio]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

15

Example 135: 6-[[2-(acetylamino)ethyl]thio]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

Example 163: 4-[(1-methyl-1H-indazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

20

Example 167: 4-[[4-hydroxy-3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 174: 4-[(3-acetylphenyl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

25

Example 184: 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

Example 185: 4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-6-(phenylsulfonyl)-3-quinolinecarboxamide,

30

Example 186: 7-methyl-4-[[3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

Example 265: 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide,

5 Example 266: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide,

Example 267: 4-[(3-acetylphenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

10

Example 268: 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

15

Example 269: 4-(2,3-dihydro-1,4-benzodioxin-5-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

Example 270: 4-[(3-chlorophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

20

Example 271: 4-[(3-cyanophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

Example 272: 4-(1,3-benzothiazol-6-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

25

Example 273: 4-[(3-fluorophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide

30

Example 285: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide

Example 287: 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide

Example 292: 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide

5 Example 294: 4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 303: 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide

10

Example 307: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 308: 8-methyl-6-(methylsulfonyl)-4-(3-pyridinylamino)-3-quinolinecarboxamide

15

Example 309: 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 311: 4-[(3-fluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

20

Example 312: 4-[(3-cyanophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

25 Example 315: 4-[(1-ethyl-1H-pyrazol-5-yl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 316: 8-methyl-4-[[5-(methyloxy)-3-pyridinyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

30

Example 317: 8-methyl-4-[(5-methyl-3-pyridinyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 369: 8-chloro-4-[(3-methylphenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 370: 8-chloro-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 371: 8-chloro-4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 372: 8-chloro-4-[(3-cyanophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 373: 8-chloro-4-[(3-fluorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 374: 8-chloro-4-[(1-methyl-1*H*-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 379: methyl 3-[(3-(aminocarbonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-quinoliny]sulfonyl]benzoate

Example 380: 6-[[3,4-bis(methyloxy)phenyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 381: 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[3,4,5-tris(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide hydrochloride

Example 382: 6-[[3,4-bis(methyloxy)phenyl]sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 383: 6-[[3-(ethyloxy)phenyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 392: 6-[[2-(acetylamino)ethyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

Example 399: 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

Example 400: 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

Example 408: 4-[(3-cyanophenyl)amino]-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide,

Example 409: 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[(1-methyl-1H-benzimidazol-6-yl)amino]-3-quinolinecarboxamide

Example 414: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide

Example 426: 6-[(1-acetyl-4-piperidiny]sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 442: 6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 443: 4-[(3-cyanophenyl)amino]-6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-3-quinolinecarboxamide

Example 445: 6-[[1-(cyclopropylcarbonyl)-4-piperidiny]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide

Example 446: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-3-quinolinecarboxamide

Example 447: 6-[[1-(cyclopropylcarbonyl)-4-piperidinyl]sulfonyl]-4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-3-quinolinecarboxamide

Example 451: 6-[(1-acetyl-4-piperidinyl)sulfonyl]-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-3-quinolinecarboxamide

Example 457: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-({2-[(methylsulfonyl)amino]ethyl}sulfonyl)-3-quinolinecarboxamide

Example 459: 6-[[1-(2-furanylcarbonyl)-4-piperidinyl]sulfonyl]-8-methyl-4-[(1-methyl-1*H*-benzimidazol-6-yl)amino]-3-quinolinecarboxamide

Example 475: 6-[[3-(dimethylamino)-3-oxopropyl]sulfonyl]-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-3-quinolinecarboxamide

Example 500: 4-[(2,3-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 501: 4-[(3-chloro-2-fluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 502: 4-[(3,5-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 539: 4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 540: 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 546: 6-({5-[(dimethylamino)carbonyl]-3-pyridinyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide hydrochloride

Example 579: 8-methyl-6-[(1-methylethyl)sulfonyl]-4-(3-pyridinylamino)-3-quinolinecarboxamide

Example 580: 6-[(1,1-dimethylethyl)sulfonyl]-8-methyl-4-(3-pyridinylamino)-3-quinolinecarboxamide

Example 584: 4-[(1-ethyl-1H-pyrazol-5-yl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

Example 585: 6-[(1,1-dimethylethyl)sulfonyl]-4-[(1-ethyl-1H-pyrazol-5-yl)amino]-8-methyl-3-quinolinecarboxamide

Example 588: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 590: 4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-3-quinolinecarboxamide

Example 591: 8-ethyl-4-[[4-fluoro-3-(methoxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 592: 8-ethyl-4-[(3-fluorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 593: 4-[(3-cyanophenyl)amino]-8-ethyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 598: 8-ethyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 599: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-ethyl-6-(methylsulfonyl)-3-quinolinecarboxamide

Example 600: 8-ethyl-6-(methylsulfonyl)-4-(3-pyridinylamino)-3-quinolinecarboxamide

Example 624: 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-fluoro-6-(methylsulfonyl)-3-quinolinecarboxamide

5

Example 666: 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-(ethylsulfonyl)-3-quinolinecarboxamide

Example 667: 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-(propylsulfonyl)-3-quinolinecarboxamide

10

Example 668: 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

Example 669: 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-3-quinolinecarboxamide

15

Example 670: 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

20

Example 671: 6-(ethylsulfonyl)-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide

Example 674: 6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide

25

Example 676: 8-chloro-4-[(5-fluoro-3-pyridinyl)amino]-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

Example 677: 8-chloro-6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-3-quinolinecarboxamide

30

Example 678: 4-[(5-chloro-3-pyridinyl)amino]-6-(ethylsulfonyl)-8-methyl-3-quinolinecarboxamide

Example 679: 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-(propylsulfonyl)-3-quinolinecarboxamide

Example 680: 4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-8-methyl-3-quinolinecarboxamide

10 and pharmaceutically acceptable salts thereof.

Preferred compounds include:

8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide,

4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

8-methyl-4-[(1-methyl-1*H*-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide,

8-methyl-4-[(5-methyl-3-pyridinyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide hydrochloride

6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide,

4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-[[1-(2-furanylcarbonyl)-4-piperidinyl]sulfonyl]-8-methyl-3-quinolinecarboxamide

5 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-({2-[(methylsulfonyl) amino] ethyl}sulfonyl)-3-quinolinecarboxamide

4-[(3,5-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

10 6-({5-[(dimethylamino)carbonyl]-3-pyridinyl}sulfonyl)-8-methyl-4-[[3-(methoxy) phenyl]amino]-3-quinolinecarboxamide hydrochloride

4-[(1-ethyl-1H-pyrazol-5-yl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

15

6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide

20

8-chloro-6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-3-quinolinecarboxamide

and pharmaceutically acceptable salts thereof.

25

Salts of the compounds of the present invention are also encompassed within the scope of the invention. Because of their potential use in medicine, the salts of the compounds of formula (I) are preferably pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts can include acid or base addition salts. A pharmaceutically acceptable acid addition salt can be formed by reaction of a compound of formula (I) with a suitable inorganic or organic acid (such as hydrobromic, hydrochloric, sulfuric, nitric, phosphoric, succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, *p*-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid), optionally in a suitable solvent such as an organic solvent, to give the salt which is usually isolated for example by crystallisation and filtration. A pharmaceutically acceptable acid addition salt of a compound of formula (I)

30

can be for example a hydrobromide, hydrochloride, sulfate, nitrate, phosphate, succinate, maleate, acetate, fumarate, citrate, tartrate, benzoate, *p*-toluenesulfonate, methanesulfonate or naphthalenesulfonate salt. A pharmaceutically acceptable base addition salt can be formed by reaction of a compound of formula (I) with a suitable inorganic or organic base, optionally in a suitable solvent such as an organic solvent, to give the base addition salt which is usually isolated for example by crystallisation and filtration. Other non-pharmaceutically acceptable salts, eg. oxalates or trifluoroacetates, may be used, for example in the isolation of compounds of the invention, and are included within the scope of this invention. The invention includes within its scope all possible stoichiometric and non-stoichiometric forms of the salts of the compounds of formula (I).


Also included within the scope of the invention are all solvates, hydrates and complexes of compounds and salts of the invention.

Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit *cis-trans* isomerism). The individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

The compounds of this invention may be made by a variety of methods, including standard chemistry. Any previously defined variable will continue to have the previously defined meaning unless otherwise indicated. Illustrative general synthetic methods are set out below and then specific compounds of the invention are prepared in the working Examples.

30 **Process a**

Compounds of formula (I), wherein R^{34} , R^{19} , R^{20} , R^1 and R^2 are as defined above, may be prepared from compounds of formula II;




wherein R^{34} , R^{19} , and R^{20} are as defined above, and X represents a halogen atom, by treatment with an amine of formula R^1R^2NH , wherein R^1 and R^2 are as defined above.

5

Suitable conditions for process a) include stirring in a suitable solvent such as acetonitrile, N,N-dimethylformamide or ethanol, at a suitable temperature, such as between room temperature and the reflux temperature of the solvent, for example at 80°C, optionally in the presence of a suitable base such as N,N-diisopropylethylamine, or in the presence of an acid catalyst such as the salt of an amine base, such as pyridine hydrochloride. Alternatively, process a) may be carried out under microwave irradiation, at a suitable power such as 100-300W, for example at 150W, in a suitable solvent such as N-methyl-2-pyrrolidinone or N,N-dimethylformamide, at a suitable temperature such as 60-200°C, for example at 150°C.

15

Compounds of formula (II), wherein R^{34} , R^{19} , R^{20} and X are as defined above, may be prepared from compounds of formula (IV);




20

wherein R^{34} , R^{19} , and R^{20} are as defined above, by treatment with a suitable chlorinating agent, such as thionyl chloride, in the presence of a suitable catalyst such as N,N-dimethylformamide, followed by treatment with ammonia under suitable conditions, such as 880 ammonia at room temperature.

25


Compounds of formula (IV), wherein R^{34} , R^{19} , and R^{20} are as defined above, may be prepared from compounds of formula (V);



5

wherein R^{34} , R^{19} , and R^{20} are as defined above, by hydrolysis with a suitable base, such as aqueous sodium hydroxide, in a suitable solvent, such as ethanol, at a suitable temperature such as room temperature.


- 10 Compounds of formula (V), wherein R^{34} , R^{19} , and R^{20} are as defined above, may be prepared from compounds of formula (VI);




- 15 wherein R^{34} , R^{19} , and R^{20} are as defined above, by heating in a suitable solvent, such as diphenyl ether, at a suitable temperature such as 200-300°C, for example at 250°C. The preparation of compounds of formulae (IV), (V), and (VI) wherein R^{34} represents MeSO_2 , R^{19} represents H and R^{20} represents H have been previously described in patent application WO 02/068394 A1 (Glaxo Group Limited).

20

Compounds of formula (VI), wherein R^{34} , R^{19} , and R^{20} are as defined above, may be prepared from compounds of formula (VII), wherein R^{34} , R^{19} , and R^{20} are as defined above, and the compound of formula (VIII);




(VII)



(VIII)

Suitable conditions include heating together compounds of formulae (VII) and (VIII) in a suitable solvent such as ethanol or in the absence of solvent, at a suitable temperature, such as 60-100°C, for example at 80°C.

Compounds of formula (VII) wherein R^{34} , R^{19} , and R^{20} are as defined above may be prepared by reduction of compounds of formula (XIV), wherein R^{34} , R^{19} , and R^{20} are as defined above;




(XIV)


suitable conditions where $n = 1$ or 2 include catalytic hydrogenation with hydrogen and a suitable catalyst, such as palladium on carbon, in a suitable solvent such as acetic acid.

Suitable conditions where $n = 0$ include reduction with a reducing agent such as iron in dilute acetic acid, at a suitable temperature such as 85-90°C.

Compounds of formula (XIV), wherein R^{34} represents $R^3S(=O)_n$, R^{19} represents hydrogen or C_{1-6} alkyl, n is 0 or 2, and R^{20} is as defined above, may be prepared from compounds of formula (XV), wherein R^{19} represents hydrogen or C_{1-6} alkyl and R^{20} is as defined above and compounds of formula (XVI) wherein R^3 is defined above and $n = 0$ or 2;



(XV)



(XVI)

Suitable conditions where $n = 0$ include treatment of the compound of formula (XV) with a thiol of formula (XVI) ($n=0$) in the presence of a suitable base such as potassium carbonate, in a suitable solvent such as acetonitrile, at a suitable temperature such as room temperature. Where $n = 2$, suitable conditions include treatment of the compound of formula (XV) with the sodium salt of a sulphinic acid of formula (XVI) ($n=2$), in a suitable solvent such as dimethylacetamide, at a suitable temperature such as 30-100°C, for example at 50°C.


Alternatively, compounds of formula (XIV) where n represents 2 may be prepared from compounds of formula (XIV) where n represents 0 by oxidation with a suitable oxidising agent, such as oxone, in a suitable solvent such as a mixture of methanol and water, at a suitable temperature such as room temperature. Compounds of formula (XIV) where n represents 1 may be prepared from compounds of formula (XIV) where n represents 0 by oxidation with a suitable oxidising agent, such as ceric ammonium nitrate, in the presence of a suitable solid support such as hydrated silica gel, in a suitable solvent such as methylene chloride, at a suitable temperature such as 20-40°C, for example at room temperature.

Compounds of formula R^1R^2NH may contain amine or acid groups which are suitably protected. Examples of suitable protecting groups and the means for their removal are well known in the art, see for instance T. W. Greene and P. G. M. Wuts 'Protective Groups in Organic Synthesis' (3rd Ed., J. Wiley and Sons, 1999). Addition or removal of such protecting groups may be accomplished at any suitable stage in the synthesis of compounds of formula (I).

Compounds of formula (II) wherein R^{34} represents $R^3S(=O)_n$, n represents 0, R^{19} represents hydrogen or C_{1-6} alkyl, X represents chlorine and R^{20} is as defined above may alternatively be prepared from compounds of formula (IX), wherein X represents chlorine, Y represents iodine, Z represents hydrogen or C_{1-6} alkyl, and R^{20} is as defined above, by treatment with a trialkylstannane of formula R^3SSnW_3 , wherein W represents a C_{1-6} alkyl group such as an n -butyl group. Suitable conditions include heating in the presence of a suitable catalyst, such as a palladium catalyst, for example tetrakis(triphenylphosphine) palladium (0), in a suitable solvent such as toluene, at a suitable temperature such as between 80°C and 150°C, for example at 110°C.

Process b

Compounds of formula (I), wherein R^1 , R^2 , R^{34} , R^{19} , and R^{20} are as defined above, and n
 5 = 0, may alternatively be prepared from compounds of formula (III);




wherein R^1 , R^2 and R^{20} are as defined above, and Z represents hydrogen, C_{1-6} alkyl or
 10 halogen for example chlorine and Y represents hydrogen, chlorine, bromine or iodine, by
 treatment with a thiol of formula R^3SH , or the sodium salt thereof, R^3SNa , wherein R^3 is
 as defined above, with the proviso that at least one of Y and Z represent halogen.

Suitable conditions for process b) include heating in a suitable solvent such as toluene or
 15 *N,N*-dimethylformamide, at a suitable temperature such as 60-150°C, for example at
 110°C, in the presence of a suitable catalyst, such as a palladium catalyst, for example
 tris(dibenzylideneacetone) palladium (II), and a suitable ligand, such as a phosphine
 ligand, for example (oxydi-2,1-phenylene)bis(diphenylphosphine), and in the presence of
 a suitable base such as potassium *tert*-butoxide.

20

Alternatively, conditions for process b) include heating in a suitable solvent such as 1,3-
 dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone or dimethoxyethane, at a suitable
 temperature such as 60-150°C, for example at 85°C, optionally in the presence of a
 suitable catalyst, such as a copper catalyst, for example copper (I) iodide, and in the
 25 presence of a suitable base such as potassium phosphate or potassium carbonate and
 optionally in the presence of a suitable ligand for example *N,N*-diethylsalicylamide.

Compounds of formula (III), wherein R^1 , R^2 , R^{20} , Y and Z are as defined above, may be prepared from compounds of formula (IX), wherein R^{20} , X, Y and Z are as defined above, by treatment with an amine of formula R^1R^2NH , wherein R^1 and R^2 are as defined above;




(IX)

suitable conditions include stirring in a suitable solvent such as acetonitrile, at a suitable temperature, such as between room temperature and the reflux temperature of the solvent, for example at $80^\circ C$, optionally in the presence of a base such as *N,N*-diisopropylethylamine, or in the presence of an acid catalyst such as pyridine hydrochloride. Alternatively, preparation of compounds of formula (III) from compounds of formula (IX) may be carried out under microwave irradiation, at a suitable power such as 100-300W, for example at 150W, in a suitable solvent such as *N*-methyl-2-pyrrolidinone, at a suitable temperature such as $60-200^\circ C$ for example at $150^\circ C$.

The compounds of formula (IX) may be prepared according to the following synthetic scheme, Scheme 1, wherein R^{19} , R^{20} , Y and Z are as defined above:

SCHEME 1




Suitable conditions for the reactions of Scheme 1 are: (A) heating together compounds of formulae (X) and (VIII) in the absence of solvent, at a suitable temperature, such as 60-100°C, for example at 80°C; (B) heating compounds of formula (XI) in a suitable solvent, such as diphenyl ether, at a suitable temperature such as 200-300°C, for example at 250°C; (C) hydrolysis of compounds of formula (XII) with a suitable base, such as aqueous sodium hydroxide, in a suitable solvent, such as ethanol, at a suitable

temperature such as room temperature; (D) treatment of compounds of formula (XIII) with a suitable halogenating agent, such as a chlorinating agent, for example thionyl chloride, in the presence of a suitable catalyst such as *N,N*-dimethylformamide, followed by treatment with ammonia under suitable conditions, such as 880 ammonia at room temperature.

Preparation of the compounds of formulae (XI) and (XII) wherein Y represents iodine and Z and R²⁰ both represent hydrogen have been previously described in: *Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry*, 2002, **41B(3)**, 650-652. Preparation of the compound of formula (XIII) wherein Y represents iodine and Z and R²⁰ both represent hydrogen has been previously described in: PCT Int. Appl. (1999), WO9932450 A1.

Compounds of formula (X) are either known compounds (for example available from commercial suppliers such as Aldrich) or may be prepared by conventional means.

Compounds of formula (V), wherein R³⁴ represents R³S(=O)_n-, R¹⁹ represents hydrogen or C₁₋₆alkyl, R²⁰ is as defined above and n = 0, may be prepared by treatment of compounds of formula (XII), wherein Y and R²⁰ are as defined above and Z represents hydrogen or C₁₋₆alkyl with a thiol of formula R³SH, wherein R³ is as defined above, according to the following scheme:



Suitable conditions for preparation of compounds of formula (V) from compounds of formula (XII) and a thiol of formula R³SH include heating in a suitable solvent such as toluene, at a suitable temperature such as 60-120°C, for example at 110°C, in the

presence of a suitable catalyst, such as a palladium catalyst, for example tris(dibenzylideneacetone) palladium (II), and a suitable ligand, such as a phosphine ligand, for example (oxydi-2,1-phenylene)bis(diphenylphosphine), and in the presence of a suitable base such as potassium *tert*-butoxide.

5

Compounds of formulae R^1R^2NH and R^3SH are either known compounds (for example available from commercial suppliers such as Aldrich) or may be prepared by conventional means.

- 10 Certain compounds of formula R^3SH may be prepared from compounds of formula R^3SSR^3 . Suitable conditions include treatment with a suitable reducing agent such as a phosphine, for example triphenylphosphine, in the presence of an acid such as concentrated hydrochloric acid, in a suitable solvent such as a mixture of water and 1,4-dioxane, at a suitable temperature such as between 20°C and 100°C, for example at
- 15 40°C. Alternatively certain compounds of formula R^3SH may be prepared from compounds of formula R^3SO_2Cl . Suitable conditions include treatment with a suitable reducing agent such as a phosphine, for example triphenylphosphine, in a suitable solvent such as 1,4-dioxane, at a suitable temperature such as between 0°C and 50°C, for example at 20°C.

20

Compounds of formula R^1R^2NH may be used in the free base form, or in the form of a suitable salt, such as a hydrochloride salt. Where the free base form is commercially available, suitable salt forms may be prepared by conventional means. Similarly, where a salt form is commercially available, the free base form may be prepared by conventional

25 means.

- Compounds of formula R^3SH may contain amine or acid groups which are suitably protected. Examples of suitable protecting groups and the means for their removal are well known in the art, see for instance T. W. Greene and P. G. M. Wuts 'Protective Groups in Organic Synthesis' (3rd Ed., J. Wiley and Sons, 1999). Addition or removal of such
- 30 protecting groups may be accomplished at any suitable stage in the synthesis of compounds of formula (I).


Process c

- Compounds of formula (I) may also be prepared by a process of interconversion between
- 5 compounds of formula (I). For example, compounds of formula (I) where $n = 2$ may be prepared from compounds of formula (I) wherein $n = 0$ or 1 , by treatment with a suitable oxidising agent, such as oxone, in a suitable solvent such *N,N*-dimethylformamide or a mixture of *N,N*-dimethylformamide and anisole, at a suitable temperature such as room temperature. Compounds of formula (I) where $n = 1$ may be prepared from compounds of
- 10 formula (I) where $n = 0$ by oxidation with a suitable oxidising agent, such as oxone or ceric ammonium nitrate, in the presence of a suitable solid support such as hydrated silica gel, in a suitable solvent such as methylene chloride, at a suitable temperature such as 20-40°C, for example at room temperature.
- 15 Alternative processes of interconversion between compounds of formula (I) may include, for example oxidation, reduction, hydrolysis, alkylation, dealkylation, amide bond formation, protection, deprotection, sulphonamide formation or substitution, using methods for functional group interconversion well known to those skilled in the art.

20


Process d

- As a particular example of a process of interconversion, compounds of formula (I), wherein R^{34} represents $R^3S(=O)_n-$, R^3 represents an aryl group substituted by -
- 25 $CONR^{29}R^{30}$, R^{19} represents hydrogen or C_{1-6} alkyl, and wherein R^1 , R^2 , R^{20} , R^{29} , R^{30} and n are as defined above, may alternatively be prepared from corresponding compounds of formula (I) in which R^3 represents an aryl group substituted by -COOH, namely compounds of formula (XVII);



wherein R^{19} represents hydrogen or C_{1-6} alkyl, and R^1 , R^2 , R^{20} and n are as defined above, by coupling with a primary or secondary amine, in a suitable solvent, such as *N,N*-dimethylformamide, in the presence of a suitable amide coupling reagent, such as *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate, optionally in the presence of a suitable base, such as *N,N*-diisopropylethylamine, at a suitable temperature, such as room temperature. (Step (I))

Compounds of formula (XVII), wherein R^{19} represents hydrogen or C_{1-6} alkyl, and R^1 , R^2 , R^{20} and n are as defined above, may be prepared from compounds of formula XVIII;



wherein R^{19} represents hydrogen or C_{1-6} alkyl, and R^1 , R^2 , R^{20} and n are as defined above, by hydrolysis with a suitable base, such as aqueous sodium hydroxide, in a suitable solvent, such as ethanol, at a suitable temperature such as 75°C. (Step (II))

Compounds of formula (XVIII) wherein $n = 2$ may be prepared from compounds of formula (XVIII) wherein $n = 0$, by treatment with a suitable oxidising agent, such as oxone, in a suitable solvent such as *N,N*-dimethylformamide, at a suitable temperature such as room temperature. (Step (III))

Compounds of formula (XVIII), wherein R^{19} represents hydrogen or C_{1-6} alkyl, R^1 , R^2 and R^{20} are as defined above, and $n=0$, may be prepared from compounds of formula (III) wherein Z represents hydrogen or C_{1-6} alkyl by treatment with a suitable thiol such as methyl 3-mercaptobenzoate or methyl 4-mercaptobenzoate (both commercially available from Toronto). Suitable conditions for this include heating in a suitable solvent such as 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone, at a suitable temperature such as 60-150°C, for example at 85°C, in the presence of a suitable catalyst, such as a copper catalyst, for example copper (I) iodide, and in the presence of a suitable base such as potassium phosphate or potassium carbonate, optionally in the presence of a suitable ligand for example *N,N*-diethylsalicylamide. (Step (IV))

The order of the steps comprising this process may be arranged in a number of different ways. For example the order of steps (II) and (III) may be reversed so that compounds of formula (I) may be prepared by step (IV) followed by step (II) followed by step (III) followed by step (I).


By a similar process, compounds of formula (I), wherein R^{34} represents $R^3S(=O)_n-$ and R^3 represents a C_{1-6} alkyl group substituted by $-CONR^{26}R^{27}$ and wherein R^{19} represents hydrogen or C_{1-6} alkyl, and R^1 , R^2 , R^{20} , R^{26} , R^{27} and n are as defined above, may alternatively be prepared from compounds of formula (III) where Z represents hydrogen or C_{1-6} alkyl and a suitable thiol such as ethyl 3-mercaptopropionate (commercially available from Aldrich) as shown in the scheme below:

Steps (I) to (IV) of Scheme 2 use conditions as described in process d above.

The order of the steps comprising this process may be arranged in a number of different ways. For example the order of steps may be changed so that compounds of formula (I) may be prepared from compounds of formula (III) by step (IV) followed by step (II) followed by step (III) followed by step (I).

Alternatively the order of steps may be changed so that compounds of formula (I) may be prepared by step (IV) followed by step (II) followed by step (I) followed by step (III).

Scheme 2




5

wherein R^3 is $\text{R}^{26}\text{R}^{27}\text{NCO}(\text{CH}_2)_2-$.

Process e

As a particular example of a process of interconversion, compounds of formula (I),
 5 wherein R^{34} represents $R^3S(=O)_n$ - and R^3 represents a piperidinyl group which is substituted by a substituent selected from C_{1-6} alkyl-, C_{1-6} alkylCO-, C_{3-7} cycloalkylCO-, heteroarylCO- (optionally substituted by one or more C_{1-4} alkyl- groups), C_{1-6} alkoxyCO-, arylCO-, $R^{31}R^{32}NCO$ -, C_{1-6} alkylSO₂-, arylSO₂- or heteroarylSO₂- (optionally substituted by one or more C_{1-4} alkyl or C_{1-4} alkylCONH- groups) and wherein R^1 , R^2 , R^{20} , R^{31} , R^{32} and n
 10 are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, may alternatively be prepared from compounds of formula (XIX);




15 wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, by treatment with an electrophile, such as an acylating agent, such as an acid chloride, in a suitable solvent, such as 1,4-dioxane, in the presence of a suitable base, such as an amine base, for example triethylamine, at a suitable temperature, such as room temperature. Alternative electrophiles that may be used for this process include
 20 sulphonyl chlorides, alkyl chloroformates, alkyl halides and acid anhydrides.

Alternatively, compounds of formula (I), wherein R^{34} represents $R^3S(=O)_n$ - and R^3 represents a piperidinyl which is substituted by a substituent selected from C_{1-6} alkylCO-, C_{3-7} cycloalkylCO-, heteroarylCO- (optionally substituted by one or more C_{1-4} alkyl- groups),
 25 or arylCO-, and wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, may alternatively be prepared from compounds of formula (XIX), by coupling with a carboxylic acid, in a suitable solvent, such as *N,N*-dimethylformamide, in the presence of a suitable amide coupling reagent, such as *O*-(7-azabenzotriazol-1-yl)-

N,N,N',N'-tetramethyluronium hexafluorophosphate, optionally in the presence of a suitable base, such as *N,N*-diisopropylethylamine, at a suitable temperature, such as room temperature. (Step (I))

- 5 Compounds of formula (XIX), wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, may be prepared from compounds of formula (XX);



- 10 wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, by treatment with a suitable reagent, such as a strong acid, for example trifluoroacetic acid, at a suitable temperature, such as room temperature (Step (II)).

- 15 Compounds of formula (XX) wherein R^1 , R^2 , and R^{20} are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and $n = 2$, may be prepared from compounds of formula (XX) wherein $n = 0$, by treatment with a suitable oxidising agent, such as oxone, in a suitable solvent such as *N,N*-dimethylformamide, at a suitable temperature such as room temperature (Step (III)).

- 20 Compounds of formula (XX) wherein R^1 , R^2 , and R^{20} are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and $n = 0$, may be prepared from compounds of formula (III) wherein R^1 , R^2 , Y and R^{20} are as defined above and Z represents hydrogen or C_{1-6} alkyl, by treatment with 1,1-dimethylethyl 4-mercapto-1-piperidinecarboxylate (prepared as described in US5317025A) (Step (IV)).

25


Suitable conditions for this process include heating in a suitable solvent such as dimethylformamide, at a suitable temperature such as 60-150°C, for example at 110°C, in the presence of a suitable catalyst, such as a palladium catalyst, for example

tris(dibenzylideneacetone) palladium (II), and a suitable ligand, such as a phosphine ligand, for example (oxydi-2,1-phenylene)bis(diphenylphosphine), and in the presence of a suitable base such as potassium *tert*-butoxide.

- 5 The order of the steps comprising this process may be arranged in a number of different ways. For example the order of steps may be changed so that compounds of formula (I) may be prepared by step (IV) followed by step (II) followed by step (I) followed by step (III).
- 10 Similarly, compounds of formula (I) wherein R^{34} represents $R^3S(=O)_n-$ and R^3 represents a C_{1-6} alkyl which is substituted by $-NR^{17}R^{18}$, $-NR^{16}COR^{15}$, C_{1-6} alkoxyCONR²⁵- or C_{1-6} alkylSO₂NR³³- and wherein R^1 , R^2 , R^{20} , R^{15} , R^{17} , R^{18} and n are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and R^{16} , R^{25} and R^{33} represent hydrogen may alternatively be prepared from compounds of formula (III), wherein R^1 , R^2 , Y and R^{20} are
- 15 as defined above and Z represents hydrogen or C_{1-6} alkyl, and a thiol such as *tert*-butyl-*N*-(2-mercaptoethyl)carbamate (Aldrich), as is illustrated in the following Scheme (Scheme 3):

Steps (I) to (IV) of Scheme 3 use conditions as described in process e above.


SCHEME 3



wherein R^{34} represents $\text{C}_{1-6}\text{alkylSO}_2-$, wherein the $\text{C}_{1-6}\text{alkyl}$ group is substituted by $\text{R}^{15}\text{CONR}^{16}-$, $\text{C}_{1-6}\text{alkoxyCONR}^{25}-$, $\text{C}_{1-6}\text{alkylSO}_2\text{NR}^{33}-$ or $\text{R}^{17}\text{R}^{18}\text{N}-$.

Process f

As a particular example of a process of interconversion compounds of formula (I),
 5 wherein R^{34} represents $R^3S(=O)_n$ - and R^3 represents an aryl group substituted by a C_{1-6} alkoxy-, C_{3-7} cycloalkoxy- or C_{3-7} cycloalkyl(C_{1-6} alkoxy)- group, R^1 , R^2 , R^{20} and n are as defined above, and R^{19} represents hydrogen or C_{1-6} alkyl, may alternatively be prepared from compounds of formula (XXI);



wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, by coupling with a suitable alkylating agent, in a suitable solvent, such as acetonitrile, in the presence of a suitable base, such as potassium carbonate, at a suitable temperature,
 15 such as 0 to 100°C, for example the reflux temperature of the solvent.

Alternatively compounds of formula (I) wherein R^{34} represents $R^3S(=O)_n$ - and R^3 represents an aryl group substituted by a C_{1-6} alkoxy-, C_{3-7} cycloalkoxy- or C_{3-7} cycloalkyl(C_{1-6} alkoxy)- group, R^1 , R^2 , R^{20} and n are as defined above, and R^{19} represents hydrogen or
 20 C_{1-6} alkyl, may be prepared from compounds of formula (XXI) by coupling with a suitable alcohol in a suitable solvent such as tetrahydrofuran, at a suitable temperature such as room temperature in the presence of a suitable coupling agent such as di-*tert* butylazodicarboxylate.

Compounds of formula (XXI) wherein $n = 2$ may be prepared from compounds of formula (XXI) wherein $n = 0$, by treatment with a suitable oxidising agent, such as oxone, in a suitable solvent such as *N,N*-dimethylformamide, at a suitable temperature such as room temperature.

- Compounds of formula (XXI) wherein R^1 , R^2 , and R^{20} are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and $n = 0$, may be prepared from compounds of formula (III) wherein R^1 , R^2 , R^{20} and Y are as defined above, and wherein Z represents hydrogen or C_{1-6} alkyl, by treatment with 4-[[*tert*-butyl(dimethyl)silyl]oxy]benzenethiol (prepared according to EP 465802 A1). Suitable conditions for this process include heating in a suitable solvent such as dimethylformamide, at a suitable temperature such as 60-150°C, for example at 110°C, in the presence of a suitable catalyst, such as a palladium catalyst, for example tris(dibenzylideneacetone) palladium (II), and a suitable ligand, such as a phosphine ligand, for example (oxydi-2,1-phenylene)bis(diphenylphosphine), and in the presence of a suitable base such as potassium *tert*-butoxide, followed by deprotection with a suitable fluoride source such as tetrabutylammonium fluoride in a suitable solvent such as tetrahydrofuran at a suitable temperature such as room temperature.
- The order of the steps comprising this process may be arranged in a number of different ways.

Process g

20

Compounds of formula (I) may also be prepared by a process of deprotection of protected derivatives of compounds of formula (I). Examples of suitable protecting groups and the means for their removal are well known in the art, see for instance T. W. Greene and P. G. M. Wuts 'Protective Groups in Organic Synthesis' (3rd Ed., J. Wiley and Sons, 1999).


25

As an example of this, compounds of formula (I) containing a primary or secondary amine group may be prepared from compounds of formula (I) where that amine group is protected, such as a carbamate group, for example as a *tert*-butyl carbamate, by deprotecting under appropriate conditions, such as treating with a strong acid, for example trifluoroacetic acid.

30

Process h

As a particular example of a process of interconversion, compounds of formula (I), wherein R^{34} represents $R^3S(=O)_n-$ and R^3 represents C_{1-6} alkoxyethyl-, R^1 , R^2 , R^{20} and n are as defined above, and R^{19} represents hydrogen or C_{1-6} alkyl may be prepared from compounds of formula (XXVIII);



(XXVIII)

wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, by alkylation with a suitable alkylating agent, in a suitable solvent, such as *N,N*-dimethylformamide, in the presence of a suitable base, such as sodium hydride, at a suitable temperature, such as 0 to 30°C, for example at room temperature.

Alternatively compounds of formula (I) wherein R^{34} represents $R^3S(=O)_n-$ and R^3 represents C_{1-6} alkoxyethyl-, R^1 , R^2 , R^{20} and n are as defined above, and R^{19} represents hydrogen or C_{1-6} alkyl may be prepared from compounds of formula (XXVIII) wherein R^1 , R^2 , R^{20} and n are as defined above and R^{19} represents hydrogen or C_{1-6} alkyl, by coupling with a suitable alcohol in a suitable solvent such as tetrahydrofuran, at a suitable temperature such as room temperature in the presence of a suitable coupling agent such as di-*tert* butylazodicarboxylate.


Compounds of formula (XXVIII) wherein R^1 , R^2 , and R^{20} are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and $n = 2$ may be prepared from compounds of formula (XXVIII) wherein $n = 0$, by treatment with a suitable oxidising agent, such as oxone, in a suitable solvent such as *N,N*-dimethylformamide, at a suitable temperature such as room temperature.

Compounds of formula (XXVIII) wherein R^1 , R^2 and R^{20} are as defined above, R^{19} represents hydrogen or C_{1-6} alkyl, and n is 0, may be prepared from compounds of formula (III) wherein R^1 , R^2 , R^{20} and Y are as defined above and Z represents hydrogen or C_{1-6} alkyl, by treatment with 2-mercaptoethanol (available from Aldrich). Suitable conditions for this process include heating in a suitable solvent such as *N,N*-dimethylformamide, at a suitable temperature such as 60-150°C, for example at 110°C, in the presence of a suitable catalyst, such as a palladium catalyst, for example tris(dibenzylideneacetone) palladium (II), and a suitable ligand, such as a phosphine ligand, for example (oxydi-2,1-phenylene)bis(diphenylphosphine), and in the presence of a suitable base such as potassium *tert*-butoxide.

The order of the steps comprising this process may be arranged in a number of different ways.

Process i

Compounds of formula (I) wherein R^{34} represents hydrogen, R^{19} represents $R^3S(=O)_n$ -, and R^1 , R^2 , R^{20} and n are as defined above, may be prepared from compounds of formula (XXIX) wherein Y represents chlorine, bromine or iodine, in particular iodine, $n = 1$ or 2, and R^1 , R^2 , R^3 and R^{20} are as defined above, by hydrogenation using a suitable hydrogenation process such as palladium on carbon in a suitable solvent such as ethanol.



Compounds of formula (XXIX) wherein Y represents chlorine, bromine or iodine, in particular iodine, $n = 1$ or 2, and R^1 , R^2 , R^3 and R^{20} are as defined above, may be

prepared from compounds of formula (XXIX) wherein $n=0$ by treatment with a suitable oxidising agent such as oxone in a suitable solvent such as *N,N*-dimethylformamide or a mixture of *N,N*-dimethylformamide and anisole at a suitable temperature such as room temperature.

5

Compounds of formula (XXIX) wherein Y represents chlorine, bromine or iodine, in particular iodine, R^1 , R^2 , R^3 and R^{20} are as defined above, and $n=0$ may be prepared from compounds of formula (III) wherein R^1 , R^2 and R^{20} are as defined above, Y represents chlorine, bromine or iodine, especially iodine, and Z represents chlorine, bromine or iodine, especially chlorine, and a thiol of formula R^3SH by heating in a suitable solvent such as 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone, at a suitable temperature such as 60-150°C for example at 100°C in the presence of a suitable base such as potassium carbonate.


10

15

Process j

As a particular example of a process of interconversion, compounds of formula (I), wherein R^{34} represents $R^3S(=O)_n$ and wherein R^3 represents:

20



and wherein R^1 , R^2 , R^{20} , and n are as defined above, and R^{19} represents hydrogen or C_{1-6} alkyl, may be prepared from compounds of formula (XXVII) in scheme 3, wherein R^1 , R^2 , and R^{20} are as defined above and Z represents hydrogen or C_{1-6} alkyl, by treatment with a suitable alkylating agent such as ethyl 4-bromobutyrate in a suitable solvent such as 1,4-dioxane at a suitable temperature such as 120°C.

25

30

The present invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use as an active therapeutic substance in a mammal such as a human. The compound or salt can be for use in the treatment and/or prophylaxis of any

of the conditions described herein and/or for use as a phosphodiesterase inhibitor, for example for use as a phosphodiesterase 4 (PDE4) inhibitor. "Therapy" may include treatment and/or prophylaxis.

- 5 Also provided is the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament (e.g. pharmaceutical composition) for the treatment and/or prophylaxis of an inflammatory and/or allergic disease in a mammal such as a human.
- 10 Also provided is a method of treatment and/or prophylaxis of an inflammatory and/or allergic disease in a mammal (e.g. human) in need thereof, which comprises administering to the mammal (e.g. human) a therapeutically effective amount of a compound of formula (I) as herein defined or a pharmaceutically acceptable salt thereof.
- 15 Phosphodiesterase 4 inhibitors are believed to be useful in the treatment and/or prophylaxis of a variety of diseases, especially inflammatory and/or allergic diseases, in mammals such as humans, for example: asthma, chronic bronchitis, emphysema, atopic dermatitis, urticaria, allergic rhinitis (seasonal or perennial), vasomotor rhinitis, nasal polyps, allergic conjunctivitis, vernal conjunctivitis, occupational conjunctivitis, infective
- 20 conjunctivitis, eosinophilic syndromes, eosinophilic granuloma, psoriasis, rheumatoid arthritis, chronic obstructive pulmonary disease (COPD) including chronic bronchitis and emphysema, septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, chronic glomerulonephritis, endotoxic shock, adult respiratory distress syndrome, multiple sclerosis or memory impairment (including Alzheimer's
- 25 disease).

In the treatment and/or prophylaxis, the inflammatory and/or allergic disease is preferably chronic obstructive pulmonary disease (COPD) including chronic bronchitis and emphysema, asthma, rheumatoid arthritis, psoriasis or allergic rhinitis in a mammal (e.g.

30 human). More preferably, the treatment and/or prophylaxis is of COPD including chronic bronchitis and emphysema or asthma in a mammal (e.g. human). PDE4 inhibitors are thought to be effective in the treatment of asthma (e.g. see M.A.Giembycz, *Drugs*, Feb. 2000, **59**(2), 193-212; Z. Huang et al., *Current Opinion in Chemical Biology*, 2001, **5**,

432-438; and refs cited therein) and COPD (e.g. see S.L. Wolda, *Emerging Drugs*, 2000, **5(3)**, 309-319; Z. Huang et al., *Current Opinion in Chemical Biology*, 2001, **5**, 432-438; and refs cited therein). COPD is often characterised by the presence of airflow obstruction due to chronic bronchitis and/or emphysema (S.L. Wolda, *Emerging Drugs*,
5 2000, **5(3)**, 309-319).

For use in medicine, the compounds of the present invention are usually administered as a pharmaceutical composition.

- 10 The present invention therefore provides in a further aspect a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable carriers and/or excipients.

The pharmaceutical composition can be for use in the treatment and/or prophylaxis of any
15 of the conditions described herein.

The compounds of formula (I) and/or the pharmaceutical composition may be administered, for example, by oral, parenteral (e.g. intravenous, subcutaneous, or intramuscular), inhaled, nasal, transdermal or rectal administration, or as topical
20 treatments (e.g. ointments or gels). Accordingly, the pharmaceutical composition is preferably suitable for oral, parenteral (e.g. intravenous, subcutaneous or intramuscular), inhaled or nasal administration. More preferably, the pharmaceutical composition is suitable for inhaled or oral administration, e.g. to a mammal such as a human. Inhaled administration involves topical administration to the lung, e.g. by aerosol or dry powder
25 composition.

A pharmaceutical composition suitable for oral administration can be liquid or solid; for example it can be a solution, a syrup, a suspension or emulsion, a tablet, a capsule or a lozenge.

30

A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable pharmaceutically acceptable liquid carrier(s), for example an aqueous solvent such as water, ethanol or glycerine, or an oil,

or a non-aqueous solvent, such as a surfactant, such as polyethylene glycol. The formulation may also contain a suspending agent, preservative, flavouring and/or colouring agent.

- 5 A pharmaceutical composition suitable for oral administration being a tablet can comprise one or more pharmaceutically acceptable carriers and/or excipients suitable for preparing tablet formulations. Examples of such carriers include lactose and cellulose. The tablet can also or instead contain one or more pharmaceutically acceptable excipients, for example binding agents, lubricants such as magnesium stearate, and/or tablet
10 disintegrants.

- A pharmaceutical composition suitable for oral administration being a capsule can be prepared using encapsulation procedures. For example, pellets containing the active ingredient can be prepared using a suitable pharmaceutically acceptable carrier and then
15 filled into a hard gelatin capsule. Alternatively, a dispersion, suspension or solution can be prepared using any suitable pharmaceutically acceptable carrier, for example an aqueous solution, aqueous gum or an oil and the dispersion, suspension or solution then filled into a hard or soft gelatin capsule.

- 20 The compounds of formula (I) and/or the pharmaceutical composition may be administered by a controlled or sustained release formulation as described in WO 00/50011.

- A parenteral composition can comprise a solution or suspension of the compound or
25 pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil. Alternatively, the solution can be lyophilised; the lyophilised parenteral pharmaceutical composition can be reconstituted with a suitable solvent just prior to administration.

- 30 Compositions for nasal or inhaled administration may conveniently be formulated as aerosols, solutions, suspensions, drops, gels or dry powders.

For compositions suitable and/or adapted for inhaled administration, it is preferred that the compound or salt of formula (I) is in a particle-size-reduced form, and more preferably the size-reduced form is obtained or obtainable by micronisation. The preferable particle size of the size-reduced (e.g. micronised) compound or salt is defined by a D50 value of about 0.5 to about 10 microns (for example as measured using laser diffraction).

Aerosol formulations, e.g. for inhaled administration, can comprise a solution or fine suspension of the active substance in a pharmaceutically acceptable aqueous or non-aqueous solvent. Aerosol formulations can be presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device or inhaler. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve (metered dose inhaler) which is intended for disposal once the contents of the container have been exhausted.

Where the dosage form comprises an aerosol dispenser, it preferably contains a suitable propellant under pressure such as compressed air, carbon dioxide or an organic propellant such as a chlorofluorocarbon (CFC) or hydrofluorocarbon (HFC). Suitable CFC propellants include dichlorodifluoromethane, trichlorofluoromethane and dichlorotetrafluoroethane. Suitable HFC propellants include 1,1,1,2,3,3,3-heptafluoropropane and 1,1,1,2-tetrafluoroethane. The aerosol dosage forms can also take the form of a pump-atomiser.

Optionally, in particular for dry powder inhalable compositions, a pharmaceutical composition for inhaled administration can be incorporated into a plurality of sealed dose containers (e.g. containing the dry powder composition) mounted longitudinally in a strip or ribbon inside a suitable inhalation device. The container is rupturable or peel-openable on demand and the dose of e.g. the dry powder composition can be administered by inhalation via the device such as the DISKUSTM device, marketed by GlaxoSmithKline. The DISKUSTM inhalation device is for example described in GB 2242134 A, and in such a device at least one container for the pharmaceutical composition in powder form (the container or containers preferably being a plurality of sealed dose containers mounted longitudinally in a strip or ribbon) is defined between two members peelably

secured to one another; the device comprises: a means of defining an opening station for the said container or containers; a means for peeling the members apart at the opening station to open the container; and an outlet, communicating with the opened container, through which a user can inhale the pharmaceutical composition in powder form from the
5 opened container.

In the pharmaceutical composition, each dosage unit for oral or parenteral administration preferably contains from 0.01 to 3000 mg, more preferably 0.5 to 1000 mg, of a compound of the formula (I) or a pharmaceutically acceptable salt thereof, calculated as
10 the free base. Each dosage unit for nasal or inhaled administration preferably contains from 0.001 to 50 mg, more preferably 0.01 to 5 mg, of a compound of the formula (I) or a pharmaceutically acceptable salt thereof, calculated as the free base.

The pharmaceutically acceptable compounds or salts of the invention can be
15 administered in a daily dose (for an adult patient) of, for example, an oral or parenteral dose of 0.01 mg to 3000 mg per day or 0.5 to 1000 mg per day, or a nasal or inhaled dose of 0.001 to 50 mg per day or 0.01 to 5 mg per day, of the compound of the formula (I) or a pharmaceutically acceptable salt thereof, calculated as the free base.

20 The compounds, salts and/or pharmaceutical compositions according to the invention may also be used in combination with one or more other therapeutically active agents, for example, a β_2 adrenoreceptor agonist, an anti-histamine, an anti-allergic agent, an anti-inflammatory agent (including a steroid), an anticholinergic agent or an antiinfective agent (e.g. antibiotics or antivirals).

25 The invention thus provides, in a further aspect, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof with one or more other therapeutically active agents, for example, a β_2 -adrenoreceptor agonist, an anti-histamine, an anti-allergic agent, an anti-inflammatory agent (including a steroid), an
30 anticholinergic agent or an antiinfective agent (e.g. antibiotics or antivirals).

Examples of β_2 -adrenoreceptor agonists include salmeterol (e.g. as racemate or a single enantiomer such as the R-enantiomer), salbutamol, formoterol, salmefamol, fenoterol or

terbutaline and salts thereof, for example the xinafoate salt of salmeterol, the sulphate salt or free base of salbutamol or the fumarate salt of formoterol. Long-acting β_2 -adrenoreceptor agonists are preferred, especially those having a therapeutic effect over a 24 hour period, such as salmeterol or formoterol.

5

Examples of anti-histamines include methapyrilene or loratadine.

Examples of anti-inflammatory steroids include fluticasone propionate and budesonide.

10 Examples of anticholinergic compounds which may be used in combination with a compound of formula (I) or a pharmaceutically acceptable salt thereof are described in WO 03/011274 A2 and WO 02/069945 A2 / US 2002/0193393 A1 and US 2002/052312 A1. For example, anticholinergic agents include muscarinic M3 antagonists, such as ipratropium bromide, oxitropium bromide or tiotropium bromide.

15

Other suitable combinations include, for example, combinations comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with other anti-inflammatory agents such as an anti-inflammatory corticosteroid; or a non-steroidal anti-inflammatory drug (NSAID) such as a leukotriene antagonist (e.g. montelukast), an iNOS
20 inhibitor, a tryptase inhibitor, an elastase inhibitor, a beta-2 integrin antagonist, an adenosine a2a agonist, a chemokine antagonist such as a CCR3 antagonist, or a 5-lipoxygenase inhibitor; or an antiinfective agent (e.g. an antibiotic or an antiviral). An iNOS inhibitor is preferably for oral administration. Suitable iNOS inhibitors (inducible nitric oxide synthase inhibitors) include those disclosed in WO 93/13055, WO 98/30537,
25 WO 02/50021, WO 95/34534 and WO 99/62875. Suitable CCR3 inhibitors include those disclosed in WO 02/26722.

The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical composition and thus a pharmaceutical composition comprising a
30 combination as defined above together with one or more pharmaceutically acceptable carriers and/or excipients represent a further aspect of the invention.

The individual compounds of such combinations may be administered either sequentially or simultaneously in separate or in combined pharmaceutical compositions.

5 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

10 The various aspects of the invention will now be described by reference to the following Biological Test Methods and Examples. These Examples are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

Biological Test Methods

PDE3, PDE4B, PDE4D, PDE5 and PDE6 Primary Assay Methods

5

The activity of the compounds can be measured as described below. Preferred compounds of the invention are selective PDE4 inhibitors, *i.e.* they inhibit PDE4 (e.g. PDE4B and/or PDE4D) more strongly than they inhibit other PDE's such as PDE3 and/or PDE5.

10

1. PDE enzyme sources and literature references.

15

Human recombinant PDE4B, in particular the 2B splice variant thereof (HSPDE4B2B), is disclosed in WO 94/20079 and also in M.M. McLaughlin *et al.*, "A low *K_m*, rolipram-sensitive, cAMP-specific phosphodiesterase from human brain: cloning and expression of cDNA, biochemical characterisation of recombinant protein, and tissue distribution of mRNA", *J. Biol. Chem.*, 1993, **268**, 6470-6476. For example, in Example 1 of WO 94/20079, human recombinant PDE4B is described as being expressed in the PDE-deficient yeast *Saccharomyces cerevisiae* strain GL62, e.g. after induction by addition of 150 μ M CuSO₄, and 100,000 x g supernatant fractions of yeast cell lysates are described for use in the harvesting of PDE4B enzyme.

20

Human recombinant PDE4D (HSPDE4D3A) is disclosed in P. A. Baecker *et al.*, "Isolation of a cDNA encoding a human rolipram-sensitive cyclic AMP phosphodiesterase (PDE IV_D)", *Gene*, 1994, **138**, 253-256.

25

Human recombinant PDE5 is disclosed in K. Loughney *et al.*, "Isolation and characterisation of cDNAs encoding PDE5A, a human cGMP-binding, cGMP-specific 3',5'-cyclic nucleotide phosphodiesterase", *Gene*, 1998, **216**, 139-147.

30

PDE3 was purified from bovine aorta. Its presence in the tissue was reported by H. Coste and P. Grondin in "Characterisation of a novel potent and specific inhibitor of type V phosphodiesterase", *Biochem. Pharmacol.*, 1995, **50**, 1577-1585.

PDE6 was purified from bovine retina. Its presence in this tissue was reported by: P. Catty and P. Deterre in "Activation and solubilization of the retinal cGMP-specific phosphodiesterase by limited proteolysis", *Eur. J. Biochem.*, 1991, **199**, 263-269; A. Tar *et al.* in "Purification of bovine retinal cGMP phosphodiesterase", *Methods in Enzymology*, 1994, **238**, 3-12; and/or D. Srivastava *et al.* in "Effects of magnesium on cyclic GMP hydrolysis by the bovine retinal rod cyclic GMP phosphodiesterase", *Biochem. J.*, 1995, **308**, 653-658.

2. Inhibition of PDE3, PDE4B, PDE4D, PDE5 or PDE6 Activity: Radioactive Scintillation Proximity Assay (SPA)

The ability of compounds to inhibit catalytic activity at PDE4B or 4D (human recombinant), PDE3 (from bovine aorta), PDE5 (human recombinant) or PDE 6 (from bovine retina) was determined by Scintillation Proximity Assay (SPA) in 96-well format. Test compounds (preferably as a solution in DMSO, e.g. 2 microlitre (μ l) volume) were preincubated at ambient temperature in Wallac Isoplates (code 1450-514) with PDE enzyme in 50mM Tris-HCl buffer pH 7.5, 8.3mM $MgCl_2$, 1.7mM EGTA, 0.05% (w/v) bovine serum albumin for 10-30 minutes. The enzyme concentration was adjusted so that no more than 20% hydrolysis of the substrate occurred in control wells without compound, during the incubation. For the PDE3, PDE4B and PDE4D assays [$5',8\text{-}^3H$]adenosine 3',5'-cyclic phosphate (Amersham Pharmacia Biotech, code TRK.559 or Amersham Biosciences UK Ltd, Pollards Wood, Chalfont St Giles, Buckinghamshire HP8 4SP, UK) was added to give 0.05 μ Ci per well and ~10nM final concentration. For the PDE5 and PDE6 assays [8-^3H]guanosine 3',5'-cyclic phosphate (Amersham Pharmacia Biotech, code TRK.392) was added to give 0.05 μ Ci per well and ~36nM final concentration. Plates e.g. containing approx. 100 μ l volume of assay mixture were mixed on an orbital shaker for 5 minutes and incubated at ambient temperature for 1 hour. Phosphodiesterase SPA beads (Amersham Pharmacia Biotech, code RPNQ 0150) suspended in buffer were added (~1mg per well) to terminate the assay. Plates were sealed and shaken and allowed to stand at ambient temperature for 35 minutes to 1hour to allow the beads to settle. Bound radioactive product was measured using a WALLAC TRILUX 1450 Microbeta scintillation counter. For inhibition curves, 10 concentrations (e.g. 1.5nM - 30 μ M) of each compound were assayed; more potent compounds were assayed over lower concentration ranges (assay concentrations were generally between 30 μ M and 50fM). Curves were analysed using

ActivityBase and XLfit (ID Business Solutions Limited, 2 Ocean Court, Surrey Research Park, Guildford, Surrey GU2 7QB, United Kingdom). Results were expressed as pIC₅₀ values.

- 5 Alternatively, the activity of the compounds can be measured in the following Fluorescence Polarisation (FP) assay:

3. Inhibition of PDE4B or PDE4D Activity: Fluorescence Polarisation (FP) Assay

- 10 The ability of compounds to inhibit catalytic activity at PDE4B (human recombinant) and PDE4D (human recombinant) was determined by IMAP Fluorescence Polarisation (FP) assay (Molecular Devices code: R8062) in 384-well format. Test compounds (small volume, e.g. 0.5 µl, of solution in DMSO) were preincubated at ambient temperature in black 384-well microtitre plates (supplier: NUNC, code 262260) with PDE enzyme in 10mM Tris-HCl buffer pH 7.2, 10mM MgCl₂, 0.1% (w/v) bovine serum albumin. 0.05% NaN₃ for 10-30 minutes. The enzyme level was set so that reaction was linear throughout the incubation.

- 20 Fluorescein adenosine 3',5'-cyclic phosphate (Molecular Devices code: R7091) was added to give ~40nM final concentration. Plates were mixed on an orbital shaker for 10 seconds and incubated at ambient temperature for 40 minutes. IMAP binding reagent (Molecular Devices code: R7207) was added (60µl of a 1 in 400 dilution in binding buffer of the kit stock suspension) to terminate the assay. Plates were allowed to stand at ambient temperature for 1hour. The FP ratio of parallel to perpendicular light was measured using an AnalystTM plate reader (from Molecular Devices Ltd). For inhibition curves, 11 concentrations (0.5nM - 30µM) of each compound were assayed; more potent compounds were assayed over lower concentration ranges (assay concentrations were generally between 30µM and 50fM). Curves were analysed using ActivityBase and XLfit (ID Business Solutions Limited). Results were expressed as pIC₅₀ values.

- 30 For a given PDE4 inhibitor, the PDE4B (or PDE4D) inhibition values measured using the SPA and FP assays can differ slightly. However, in a regression analysis of 100 test compounds, the pIC₅₀ inhibition values measured using SPA and FP assays have been found generally to agree within 0.5 log units for PDE4B and PDE4D (linear regression coefficient 0.966 for PDE4B and 0.971 for PDE4D; David R.Mobbs *et al*, "Comparison of

the IMAP Fluorescence Polarisation Assay with the Scintillation Proximity Assay for Phosphodiesterase Activity", poster presented at 2003 Molecular Devices UK & Europe User Meeting, 2nd October 2003, Down Hall, Harlow, Essex, United Kingdom).

- 5 Examples of compounds of the invention described above inhibit the catalytic activity at the PDE4B (human recombinant) enzyme with pIC_{50} values in the range 6.0-11.7. Biological Data obtained for some of the Examples (PDE4B and PDE5 inhibitory activity) is as follows:

Example No.	PDE4B mean pIC_{50}	PDE5 mean pIC_{50}
27	8.4	4.8
70	7.3	5.0
92	7.7	<4.5
125	6.6	5.5
265	11.3	5.2
307	10.5	<4.6
309	10.1	<4.9
312	9.4	<4.5
369	9.5	5.1
380	11.4	<7.0
399	>11.6	5.6
400	11.0	<5.0
408	11.4	4.9
446	11.3	<4.5
457	11.0	<5.5
502	8.9	<5
546	10.7	4.7

10 4. Emesis:

Many known PDE4 inhibitors cause emesis and/or nausea to greater or lesser extents (e.g. see Z. Huang *et al.*, *Current Opinion in Chemical Biology*, 2001, 5, 432-438, see especially pages 433-434 and references cited therein). Therefore, it would be preferable but not essential that a PDE4 inhibitory compound of the invention causes only limited or manageable emetic side-effects. Emetic side-effects can for example be measured by the emetogenic potential of the compound when administered to

ferrets; for example one can measure the time to onset, extent, frequency and/or duration of vomiting and/or writhing in ferrets after oral or parenteral administration of the compound. See for example A. Robichaud *et al.*, "Emesis induced by inhibitors of PDE IV in the ferret" *Neuropharmacology*, 1999, **38**, 289-297, erratum *Neuropharmacology*, 2001, **40**, 465-465.

5

EXAMPLES

In this section, "intermediates" represent syntheses of intermediate compounds intended for use in the synthesis of the "Examples".

5 **Abbreviations** used herein:

HPLC high performance liquid chromatography

NMR nuclear magnetic resonance

LC/MS liquid chromatography/mass spectroscopy

TLC thin layer chromatography

10 SPE solid phase extraction column. Unless otherwise specified the solid phase will be silica gel. C18 SPE refers to reverse phase SPE columns (eg Varian Bond Elut C18 columns). Aminopropyl SPE refers to a silica SPE column with aminopropyl residues immobilised on the solid phase (eg. IST Isolute™ columns). It is thought that compounds isolated by SPE are free bases.

15 SCX solid phase extraction (SPE) column with benzene sulfonic acid residues immobilised on the solid phase (eg. IST Isolute™ columns). When eluting with ammonia/ methanol, it is thought that compounds isolated by SCX are free bases.

20

General Experimental Details

LC/MS (Liquid Chromatography/Mass Spectroscopy)

Waters ZQ mass spectrometer operating in positive ion electrospray mode, mass range 100-1000 amu.

25 UV wavelength : 215-330nm

Column : 3.3cm x 4.6mm ID, 3µm ABZ+PLUS

Flow Rate : 3ml/min

Injection Volume : 5µl

Solvent A : 95% acetonitrile + 0.05% formic acid

30 Solvent B : 0.1% formic acid + 10mM ammonium acetate

Gradient: Mixtures of Solvent A and Solvent B are used according to the following gradient profiles (expressed as % Solvent A in the mixture): 0% A/0.7min, 0-100% A/3.5min, 100% A/1.1min, 100-0% A/0.2min

Mass Directed Automated Preparative HPLC Column, Conditions and EluentMethod A

The preparative column used was a Supelcosil ABZplus (10cm x 2.12cm internal
5 diameter; particle size 5 μ m)

UV detection wavelength : 200-320nm

Flow rate: 20ml/min

Injection Volume: 0.5ml

Solvent A: 0.1% formic acid

10 Solvent B: 95% acetonitrile + 0.05% formic acid

Gradient systems: mixtures of Solvent A and Solvent B are used according to a choice of
5 generic gradient profiles (expressed as % Solvent B in the mixture), ranging from a start
of 0 to 50% Solvent B, with all finishing at 100% Solvent B to ensure total elution.

15 It is thought that compounds isolated by this method are free bases, unless the R¹ or R³
groups contain basic moieties, in which case formate salts may be formed.

Mass Directed Automated Preparative HPLC column, conditions and eluentMethod B

20 The preparative column used was a Supelcosil ABZplus (10cm x 2.12cm internal
diameter; particle size 5 μ m)

UV detection wavelength: 200-320nm

Flow rate: 20ml/min

Injection Volume: 0.5ml

Solvent A: water + 0.1% trifluoroacetic acid

25 Solvent B: acetonitrile + 0.1% trifluoroacetic acid

Gradient systems: mixtures of Solvent A and Solvent B are used according to a choice of
5 generic gradient profiles (expressed as % Solvent B in the mixture), ranging from a start
of 0 to 50% Solvent B, with all finishing at 100% Solvent B to ensure total elution.

30 It is thought that compounds isolated by this method are trifluoroacetate salts.

Mass Directed Automated Preparative HPLC column, conditions and eluentMethod C

This is identical to method A. After purification but before solvent removal an excess (between a few drops and 0.5ml) of dilute hydrochloric acid is added to the product containing fractions.

It is thought that compounds isolated by this method are hydrochloride salts.

5

Product isolation by filtration directly from the reaction mixture

It is thought that compounds isolated by this method from reactions involving displacement of a 4-chloroquinoline intermediate with an amine of formula R^1R^2NH are hydrochloride salts.

10

'Hydrophobic Frit'

This refers to a Whatman PTFE filter medium (frit), pore size 5.0 μ m, housed in a polypropylene tube.

15

Oasis cartridge

This refers to a Waters OasisTM HLB Liquid Phase Extraction Cartridge

Evaporation of product fractions after purification

20

Reference to column chromatography, SPE and preparative HPLC purification includes evaporation of the product containing fractions to dryness by an appropriate method.

Aqueous ammonia solutions

'880 Ammonia' or '0.880 ammonia' refers to concentrated aqueous ammonia (specific gravity 0.880).


25

Intermediates and Examples

All reagents not detailed in the text below are commercially available from established suppliers such as Sigma-Aldrich.

30


Intermediate 1. 1-(Cyclopentylthio)-4-nitrobenzene



Cyclopentanethiol (1.0g) (available from Aldrich) was dissolved in acetonitrile (10ml) and potassium carbonate (1.35g) was added. After 5 min, 1-fluoro-4-nitrobenzene (1.38g) (available from Aldrich) was added and the mixture was stirred at room temperature overnight. The mixture was diluted with water and extracted with ether. The organic layer was washed with 1M aqueous sodium hydroxide (20ml), water (20ml), and 1M aqueous hydrochloric acid (20ml). The organic layer was separated and the solvent evaporated *in vacuo* to give the title compound as a yellow liquid (0.7g).

¹HNMR (CDCl₃) δ 8.12 (2H,m), 7.33 (2H,m), 3.75 (1H,m), 2.19 (2H,m), 1.87-1.62 (6H,m).


Intermediate 2. 1-(Cyclopentylsulfonyl)-4-nitrobenzene



Intermediate 1 (0.7g) was dissolved in methanol (20ml) and the solution cooled to 0°C. A solution of oxone (1.93g) in water (20ml) was added and the mixture was stirred under nitrogen for 2h at room temperature. The mixture was extracted with dichloromethane, the layers were separated (hydrophobic frit), and the organic layer evaporated to give the title compound as a yellow oil which crystallised on standing (0.79g).

LC/MS R_t 3.05 min, *m/z* 273 [MNH₄⁺]


Intermediate 3. 4-(Cyclopentylsulfonyl)aniline



Intermediate 2 (13.1g) was dissolved in acetic acid (150ml) and hydrogenated over palladium on activated carbon (1.6g) with stirring overnight. The mixture was filtered through Celite filter aid, and the filtrate was evaporated to give a yellow/green oil. The oil was taken up in methanol and an insoluble precipitate filtered off; the filtrate was evaporated *in vacuo* to give a yellow solid. Trituration with ether and filtration gave the title compound as a pale yellow solid (8.1g).

LC/MS R_t 2.5 min, m/z 243 $[MNH_4^+]$

Intermediate 4. Diethyl ({[4-cyclopentylsulfonyl]phenyl}amino)methylidene)propanedioate




Intermediate 3 (10.8g) (*Helvetica Chimica Acta* 1983, **66**(4), 1046-52) and diethyl (ethoxymethylene)malonate (11.4g) (available from Aldrich) were heated at 130°C for 2h. After cooling, the brown oil was scratched around the edge of the flask which caused the oil to solidify. Trituration with methanol gave a beige solid, which was filtered off to give the title compound (12.3g). The filtrate was evaporated *in vacuo* to give a brown oil. Purification by chromatography on silica gel, eluting with 5% ethyl acetate/cyclohexane

followed by 10% ethyl acetate/cyclohexane, gave an orange solid; trituration with methanol and filtration gave the title compound as a yellow solid (2.5g; total yield 14.8g).
LC/MS R_t 3.27min m/z 396 $[MH^+]$

5

Intermediate 5. Ethyl 6-(cyclopentylsulfonyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylate




10 Intermediate 4 (14.7g) was dissolved in diphenyl ether (150ml) and the solution heated at 250°C for 30min. After cooling, the mixture was diluted with cyclohexane and the resulting precipitate filtered off and washed with further cyclohexane to give the title compound as a beige solid (10.9g).


LC/MS R_t 2.46min m/z 350 $[MH^+]$

15

Intermediate 6. 6-(Cyclopentylsulfonyl)-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid



20 Intermediate 5 (10.9g) was dissolved in ethanol (100ml) and 2M sodium hydroxide (100ml), and the mixture was heated under reflux for 3h. After cooling, the solvent was evaporated *in vacuo* and the residue was dissolved in water and washed with ethyl acetate. The aqueous layer was acidified with 2M hydrochloric acid to between pH 5 and pH 6 which caused a precipitate to form. The precipitate was filtered off, washed with
25 water, and dried *in vacuo* overnight to give the title compound as a beige solid (9.47g).

LC/MS R_t 2.65min m/z 322 $[MH^+]$ **Intermediate 7. 4-Chloro-6-(cyclopentylsulfonyl)-3-quinolinecarboxamide**

5


Intermediate 6 (1.43g) was suspended in thionyl chloride (20ml) and *N,N*-dimethylformamide (5 drops) was added. The mixture was heated under reflux for 2h. After cooling, the thionyl chloride was evaporated *in vacuo* and the resulting residue was azeotroped with toluene. 0.880 Ammonia (25ml) was added dropwise to the yellow solid (caution: exotherm), and the suspension was stirred at room temperature for 16h. The resulting precipitate was filtered off, washed with water, and dried *in vacuo* to give the title compound (0.71g).

10

LC/MS R_t 2.47min m/z 339 $[MH^+]$

15


Similarly prepared were the following:



Intermediate No.	R^3 -	R^{19} -	R^{20} -	Starting material/ source	LCMS MH^+	LCMS R_t (min)
Intermediate 8	Ph-	H-	H-	4-(phenylsulfonyl)aniline / Maybridge	347	2.58
Intermediate 9	Me-	H-	H-	4-(methylsulfonyl)aniline / Salor	285	2.00

Intermediate 16	Ph-	H-	Me-	Intermediate 15	361	2.78
Intermediate 17	Me-	Me-	H-	1-fluoro-2-methyl-4-nitrobenzene / Aldrich	299	2.19
Intermediate 95	^t Bu-	H-	H-	4-[(1,1-dimethylethyl)sulphonyl]aniline / Helvetica Chimica Acta (1983), 66(4), 1046-52	327	2.40

Intermediate 30. 4-Chloro-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide




5

This was made in the same manner as Intermediate 7 starting from 4-[(1-methylethyl)sulfonyl]aniline (*Helvetica Chimica Acta* (1983), **66**(4), 1046-52).
LCMS R_t 2.27min m/z 313 $[MH^+]$


10

The following were also made in the same manner as Intermediate 7, with the proviso that the intermediates of formula




15

were prepared from the appropriate 4-fluoronitrobenzene in a similar manner to Intermediate 15:



Intermediate Number	R ³ SO ₂ -	R ²⁰ -	Starting Nitroaryl Compound / Supplier	Starting Sulfinic Acid / Supplier	LCMS MH ⁺	LCMS R _t (min)
Intermediate 31		H-	4-fluoronitrobenzene/ Aldrich	Sodium 4-(methoxy) benzenesulfinate / WO 9830566 A1	379	2.72
Intermediate 32		Me-	5-fluoro-2-nitrotoluene/ Aldrich	Sodium 4-(methoxy) benzenesulfinate / WO 9830566 A1	391	2.91
Intermediate 34		Me-	5-fluoro-2-nitrotoluene/ Aldrich	Sodium 4-methylbenzene sulfinate/ Aldrich	375	2.93
Intermediate 33	MeSO ₂ -	Me-	5-fluoro-2-nitrotoluene/ Aldrich	Methanesulfinic acid sodium salt/ Lancaster	299	2.12
Intermediate 50	MeSO ₂ -	MeO-	4-fluoro-2-(methoxy)-1-nitrobenzene/ Maybridge	Methanesulfinic acid sodium salt/ Lancaster	315	1.99

5 **Intermediate 10.** Diethyl {[(4-iodophenyl)amino]methylidene}propanedioate




A mixture of 4-iodoaniline (208g) (available from Aldrich) and diethyl (ethoxymethylene)malonate (210ml) (available from Aldrich) was heated to ca. 60°C, whereupon the mixture set solid. Heating was continued to 100°C, and then the mixture was removed from heating and broken up. Heating was continued at 100°C for 1h, and the solid was collected, washed with cyclohexane (1L) and ethanol (2x500ml), and dried *in vacuo* at 40°C overnight to give the title compound as a white solid (356g).

LC/MS R_t 3.57min m/z 390 $[MH^+]$

10


Intermediate 11. Ethyl 6-iodo-4-oxo-1,4-dihydro-3-quinolinecarboxylate



Diphenyl ether (175ml) was heated to reflux temperature, and Intermediate 10 was gradually added down an air condenser. Once all the reagent had been added the mixture was heated under reflux for a further 30min. The mixture was then cooled and 2-methylpentane (200ml) was added. The solid formed was collected by filtration to give the title compound (24.6g).


$^1\text{H NMR}$ (DMSO) δ 8.58 (1H,s), 8.42(1H,d), 7.99 (1H,dd), 7.44(1H,d), 4.21(2H,q), 1.28 (3H,t)

Intermediate 12. 6-Iodo-4-oxo-1,4-dihydro-3-quinolinecarboxylic acid




Sodium hydroxide (9.8g) was dissolved in water (61ml) and ethanol (30ml) was added. The resultant solution was added to Intermediate 11, and the mixture was heated under reflux for 60min with stirring under nitrogen. Concentrated hydrochloric acid was added, giving a white precipitate. After stirring for 16h, the precipitate was filtered off, washed with water and dried *in vacuo* to give the title compound as a white solid (8.15g).
LC/MS R_t 3.01min m/z 316 $[MH^+]$

10 **Intermediate 13. 4-Chloro-6-iodo-3-quinolinecarboxamide**




Intermediate 12 (8.1g) was added portionwise to stirred thionyl chloride (60ml). *N,N*-dimethylformamide (3 drops) was added and the mixture was heated under reflux with stirring under nitrogen for 1.75h. The excess thionyl chloride was evaporated *in vacuo* and the residue was azeotroped with toluene (2x50ml). The resulting pale yellow solid was added portionwise to stirred 0.880 ammonia (250ml), and the mixture stirred at room temperature for 1.5h. The solid was filtered off, washed with water and dried *in vacuo* at 60°C for 16h to give the title compound as a white solid (7.94g).
LC/MS R_t 2.72min m/z 332 $[MH^+]$

The following were made in the same manner as Intermediate 13:



Intermediate Number	R ¹⁹ -	R ²⁰ -	Starting Material	LCMS MH ⁺	LCMS R _t (min)
Intermediate 48	H-	Me-	4-iodo-2-methylaniline / Aldrich	347	3.06
Intermediate 49	H-	Cl-	2-chloro-4-iodoaniline / Avocado	367	2.99
Intermediate 72	H-	Et-	Intermediate 73	361	3.22
Intermediate 87	H-	F-	2-Fluoro-4-iodoaniline / Aldrich	352	2.65
Intermediate 67	Cl-	H-	from 3-chloro-4-iodoaniline / Aldrich	367	3.07

Intermediate 68. 4,7-Dichloro-8-methyl-3-quinolinecarboxamide




5

Intermediate 68 was prepared from 2-methyl-3-chloroaniline (Aldrich) in a similar manner to Intermediate 13.

LC/MS R_t 3.00min *m/z* 255 [MH⁺]

10 **Intermediate 14. 6-Iodo-4-([3-(methyloxy)phenyl]amino)-3-quinolinecarboxamide hydrochloride**




70

Intermediate 13 (5.0g) was dissolved in ethanol (60ml), 3-methoxyaniline (3.37ml) (available from Aldrich) was added, and the mixture was heated under reflux for 2.5h. The resulting precipitate was filtered off and washed with ether to give the title compound.

5 LC/MS R_t 2.59min m/z 420 $[MH^+]$









The following were made in the same manner as Intermediate 14, using acetonitrile as solvent:



10

Intermediate Number (a)	R¹NH-	R²⁰-	Starting Material	Amine / Source	LCMS MH⁺	LCMS R _t (min)
Intermediate 38 HCl		Cl-	Intermediate 49	3-methylaniline / Aldrich	438	3.56
Intermediate 35 HCl		Me-	Intermediate 48	4-fluoro-3-methoxyaniline / Apollo-Chem	452	2.78
Intermediate 36 HCl		Me-	Intermediate 48	2,3-dihydro-1-benzofuran-4-amine hydrobromide / <i>Journal of Heterocyclic Chemistry</i> (1980), 17 (6), 1333-5.	446	2.81
Intermediate 39 HCl		Cl-	Intermediate 49	4-fluoro-3-methoxyaniline / Apollo-Chem	472	3.29

Intermediate 40 HCl		Cl-	Intermediate 49	2,3-dihydro-1-benzofuran-4-amine hydrobromide / <i>Journal of Heterocyclic Chemistry</i> (1980), 17(6) , 1333-5.	466	3.35
Intermediate 41 HCl		Cl-	Intermediate 49	3-aminobenzonitrile / Aldrich	449	3.19
Intermediate 42 HCl		Cl-	Intermediate 49	3-fluoroaniline / Aldrich	442	3.40
Intermediate 43 HCl		Cl-	Intermediate 49	1-methyl-1H-indazol-6-amine hydrochloride / <i>Synthetic Communications</i> (1996), 26(13) , 2443-2447.	477	3.05
Intermediate 44 HCl		Me-	Intermediate 48	1-methyl-1H-benzimidazol-6-amine / <i>Heterocycles</i> (1991), 32(5) , 1003-12.	458	2.03
Intermediate 45 HCl		Me-	Intermediate 48	3-methoxyaniline / Aldrich	434	2.75
Intermediate 46 HCl		Me-	Intermediate 48	3-aminobenzonitrile / Aldrich	429	2.93
Intermediate 61 HCl		Me-	Intermediate 48	3-fluoroaniline / Aldrich	422	3.02


Intermediate 74 HCl		Et-	Intermediate 72	4-fluoro-3-methoxyaniline / Apollo-Chem	466	2.92
Intermediate 75 HCl		Et-	Intermediate 72	3-fluoroaniline / Aldrich	436	3.24
Intermediate 76 HCl		Et-	Intermediate 72	3-chloroaniline / Aldrich	452	3.44
Intermediate 77 HCl		Et-	Intermediate 72	3-aminobenzonitrile / Aldrich	443	3.12
Intermediate 78 HCl		Et-	Intermediate 72	3-methylaniline / Aldrich	432	3.15
Intermediate 79 HCl		Et-	Intermediate 72	1-methyl-1H-indazol-6-amine hydrochloride / <i>Synthetic Communications</i> (1996), 26(13) , 2443-2447	472	2.8
Intermediate 80 HCl		Et-	Intermediate 72	2,3-dihydro-1-benzofuran-4-amine hydrobromide / <i>Journal of Heterocyclic Chemistry</i> (1980), 17(6) , 1333-5.	460	2.97
Intermediate 88 HCl		F-	Intermediate 87	2,3-dihydro-1-benzofuran-4-amine hydrobromide / <i>Journal of Heterocyclic Chemistry</i> (1980),	450	3.06

				17(6), 1333-5.		
Intermediate 89 HCl		F-	Intermediate 87	3-fluoroaniline / Aldrich	426	3.11
Intermediate 90 HCl		F-	Intermediate 87	3-chloroaniline / Aldrich	442	3.19
Intermediate 91 HCl		F-	Intermediate 87	3-methylaniline / Aldrich	422	3.15
Intermediate 92 HCl		F-	Intermediate 87	3-aminobenzonitrile / Aldrich	433	2.88
Intermediate 93 HCl		F-	Intermediate 87	1-methyl-1H- indazol-6-amine hydrochloride / <i>Synthetic Communications (1996), 26(13), 2443-2447</i>	462	2.87
Intermediate 94 HCl		F-	Intermediate 87	4-fluoro-3- methoxyaniline/ Apollo-Chem	456	3.11

(a) Salt form: HCl = hydrochloride

Intermediate 63. 7-Chloro-6-iodo-4-{[3-(methoxy)phenyl]amino}-3-quinolinecarboxamide hydrochloride

5




Intermediate 63 was prepared from Intermediate 67 in a similar manner to Intermediate 14, using acetonitrile as solvent.

LC/MS R_t 3.15min m/z 452 $[MH^+]$

Intermediate 66. 7-Chloro-8-methyl-4-([3-(methoxy)phenyl]amino)-3-quinolinecarboxamide hydrochloride

5




Intermediate 66 was prepared from Intermediate 68 using 3-methoxyaniline in a similar manner to Intermediate 14.

LC/MS R_t 2.80min m/z 342 $[MH^+]$

10

Intermediate 104. 7-Chloro-4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-3-quinolinecarboxamide hydrochloride.




15 Intermediate 104 was prepared from Intermediate 68 using 2,3-dihydro-1-benzofuran-4-amine in a similar manner to Intermediate 14, using acetonitrile as solvent.

LC/MS R_t 2.80min m/z 354 $[MH^+]$

Intermediate 15. 3-Methyl-4-nitrophenyl phenyl sulphone

20




4-Fluoro-2-methyl-1-nitrobenzene (2.6g) (available from Aldrich) and sodium benzenesulfinate (3.0g) (available from Aldrich) were heated in *N,N*-dimethylacetamide (40ml) at 50°C for 16h. After cooling the mixture was filtered, the filtrate collected and the solvent removed *in vacuo*. The residue was triturated with cyclohexane and the resultant precipitate collected by filtration to give the title compound as a white solid (3.5g).

LC/MS R_f 3.22min m/z 295 $[MNH_4^+]$

10


Intermediate 18. 3-Amino-*N*-hydroxybenzenecarboximidamide



To a stirred solution of 3-aminobenzonitrile (4.0g) (available from Aldrich) in ethanol (100ml) was added hydroxylamine hydrochloride (4.7g) and potassium carbonate (14.0g) and the mixture heated under reflux for 22h. After cooling to room temperature the mixture was filtered through 'hyflo' filter aid, the residue washed with ethanol, and the filtrates concentrated *in vacuo* to give the title compound as a viscous oil (5.3g).

TLC SiO_2 (ethyl acetate) R_f = 0.34


Intermediate 19. 3-(5-Methyl-1,2,4-oxadiazol-3-yl)aniline



To a stirred solution of Intermediate 18 (5.3g) in dry tetrahydrofuran (50ml) was added 4Å molecular sieves, followed by sodium hydride (60% dispersion in mineral oil; 1.5g) and the mixture heated at 65°C for 30 min. After cooling to room temperature methyl acetate (2.8ml) was added and the mixture heated under reflux for 16h. After cooling to 20°C the mixture was added to water (100ml) and extracted with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified by chromatography on silica gel, eluting with dichloromethane to give the title compound as a white solid (4.0g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.22

Intermediate 20. 1-[2-Amino-3-chloro-6-(methyloxy)phenyl]-2-chloroethanone




Boron trichloride (25g) was added to dry dichloromethane (250ml) at 0°C and the resulting solution stirred for 10 min. A solution of 2-chloro-5-(methyloxy)aniline (30.6g) (available from Pfaltz Bauer) in dichloromethane (100ml) was added dropwise over 15 min to give a dark red/black mixture which was stirred for 20 min at 0°C. Chloroacetonitrile (29.5ml) was added, followed by the portionwise addition of aluminum chloride (28.4g). The mixture was stirred at room temperature for 1h and then heated

under reflux for 3h. The mixture was cooled in an ice/water bath and 2M hydrochloric acid added, followed by 5M hydrochloric acid (200ml). The resulting biphasic mixture was stirred at room temperature for 15h and then heated at 80°C for 30min. After cooling to room temperature the organic layer was collected and the aqueous layer extracted with
5 dichloromethane. The combined organic layers were washed with water, dried over sodium sulphate and concentrated *in vacuo* to give the title compound as a dark khaki solid (57.8g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.52

10


Intermediate 21. 4-Amino-5-chloro-1-benzofuran-3(2H)-one



15 To a stirred suspension of aluminum chloride (77.6g) in dry dichloromethane (300ml) was added dropwise a solution of Intermediate 20 (45g) in dichloromethane (250ml). The mixture was heated under reflux for 6h and then cooled to room temperature. The mixture was decomposed by the dropwise addition of 2M hydrochloric acid, then methanol and dichloromethane were added and the organic layer collected. The aqueous
20 layer was extracted with dichloromethane and the combined organic layers dried over sodium sulphate and concentrated *in vacuo*. The residue was dissolved in boiling methanol and an excess of triethylamine added. The solvent was removed *in vacuo* and the residue absorbed onto silica gel. Purification by chromatography on silica gel eluting with a gradient of 20% to 50% ethyl acetate in cyclohexane gave the title compound as an
25 orange/brown solid (46.9g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.66

Intermediate 22. N-(5-Chloro-3-oxo-2,3-dihydro-1-benzofuran-4-yl)-2,2,2-trifluoroacetamide




- 5 To a stirred solution of Intermediate 21 (2g) in dichloromethane (35ml) at 0°C was added triethylamine (2.1ml) and trifluoroacetic anhydride (2.1ml) and the mixture stirred at 0°C for 30 min, then at room temperature for 30 min. The mixture was quenched by the dropwise addition of water, the organic layer washed with water and the combined aqueous layers re-extracted with dichloromethane. The combined organic layers were
- 10 dried over sodium sulphate and concentrated *in vacuo*. Purification by chromatography on silica gel, eluting with 10% ethyl acetate in cyclohexane, gave the title compound as a bright yellow/orange solid (1.0g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.69

15

Intermediate 23. N-(5-Chloro-3-methylidene-2,3-dihydro-1-benzofuran-4-yl)-2,2,2-trifluoroacetamide




- 20 To a mixture of potassium *tert*-butoxide (2.0) and methyltriphenylphosphonium iodide (7.1g) was added dry toluene (70ml) and the mixture stirred at room temperature for 30 min, then heated under reflux for 30 min. The mixture was cooled to room temperature, a solution of Intermediate 22 (1.0g) in toluene (30ml) added dropwise and the mixture heated under reflux for 30 min. The mixture was cooled and quenched by the dropwise

addition of saturated ammonium chloride solution. The mixture was partitioned between ethyl acetate and water and the organic phase washed with water. The combined aqueous layers were re-extracted with ethyl acetate and the combined organic layers dried over sodium sulphate and concentrated *in vacuo* to give a dark brown oil.

- 5 Purification by column chromatography on silica gel, eluting with 10% ethyl acetate in cyclohexane, gave the title compound as a rose coloured solid (0.5g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.50

10 **Intermediate 24. 2,2,2-Trifluoro-N-(3-methyl-2,3-dihydro-1-benzofuran-4-yl)acetamide**




- 15 A solution of Intermediate 23 (0.10g) in ethanol (20ml) was added to 10% palladium on carbon (0.20g) and the mixture stirred under an atmosphere of hydrogen for 20h. The mixture was filtered through 'hyflo' filter aid, washed with ethanol and the solvent removed *in vacuo* to give the title compound as a white solid (0.092g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.53

20

Intermediate 25. 3-Methyl-2,3-dihydro-1-benzofuran-4-amine




25

To a stirred solution of Intermediate 24 (0.087g) in 2:2:1 tetrahydrofuran:methanol:water (5ml) was added lithium hydroxide (0.149g) and the mixture stirred at room temperature for 47h, then at 60°C for 2.5h. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate and water. The aqueous layer was re-extracted with
5 dichloromethane and the combined organic layers dried over sodium sulphate and concentrated *in vacuo* to give the title compound as a colourless oil (0.049g).

TLC SiO₂ (30% ethyl acetate in cyclohexane) R_f = 0.69


10 **Intermediate 26. 3-(1-Methyl-1H-pyrazol-3-yl)aniline**



A solution of 3-(4-bromo-1-methyl-1H-pyrazol-3-yl)aniline (1.0g) (available from
15 Maybridge) in ethanol (20ml) was added to a pre-hydrogenated suspension of 5% palladium on charcoal (0.5g) in ethanol (40ml). The resulting suspension was stirred under an atmosphere of hydrogen for 3h. The mixture was filtered through 'hyflo' filter aid and the filter pad washed with ethanol (50ml). The combined filtrate was evaporated *in vacuo* to give a brown gum. This gum was treated with 2M sodium carbonate solution
20 (100ml) and extracted with ethyl acetate (2x100ml); the organic layer was dried over magnesium sulphate and the solvent removed *in vacuo*. Purification by chromatography on silica gel, eluting with diethyl ether, gave the title compound as a white crystalline solid (0.5g).

TLC SiO₂ (diethyl ether) R_f = 0.28


25 **Intermediate 27. 1,2-Dimethyl-1H-benzimidazol-6-amine**



To a stirred solution of tin (II) chloride dihydrate (4.7g) in concentrated hydrochloric acid (15ml) was added 1,2-dimethyl-6-nitro-1*H*-benzimidazole (1g) (*J.Chem.Soc.*,**1931**, 1143-1153), and the mixture was stirred at room temperature for 6 h. The mixture was poured onto ice and chloroform, and basified to pH 10 by the addition of 10M sodium hydroxide solution. The mixture was extracted several times with chloroform and the combined organic extracts dried and concentrated *in vacuo* to give a brown solid. This was crystallised from ethanol to give the title compound.

TLC SiO₂ (dichloromethane:methanol:880 ammonia 90:10:1) R_f 0.75.


Intermediate 28. 3-Mercapto-*N,N*-dimethylbenzamide



Iodine (1g) was added to a stirred solution of 3-[(dimethylamino)carbonyl]benzenesulfonyl chloride (2g) (Borthwick *et al*, *J. Med. Chem* **2002**, 45(1), 1-18) and triphenylphosphine (8.4g) in 1,4-dioxane at 0°C. The mixture was stirred for 0.5h at ambient temperature. The mixture was poured into a sodium sulphite solution (50ml), extracted into ethyl acetate (2x30ml) and washed with 2N sodium hydroxide solution (2x40ml). The alkaline extracts were acidified and re-extracted into dichloromethane (3x50ml). The extracts were washed with water (100ml), dried (Na₂SO₄) and evaporated to give the title compound as a colourless solid (1.1g).

LC/MS R_t 2.32min, *m/z* 182 [MH⁺]

Intermediate 28. 3-Mercapto-*N,N*-dimethylbenzamide (alternative synthesis)




A solution of 3,3'-dithiobis(*N,N*-dimethylbenzamide) (Ger. Offen. (1978), DE 2821410)

5 (54.2g) in 1,4-dioxane (400ml) and water (100ml) was warmed to 35°C and concentrated hydrochloric acid (3ml) was added. Triphenylphosphine (55g) was added portionwise over 25min maintaining the temperature below 42°C, then the mixture was stirred at 40°C for 2.5h. After cooling to ambient temperature the mixture was concentrated to ca. 200ml, and partitioned between 2N aqueous sodium hydroxide solution (250ml) and ethyl acetate (500ml). The aqueous phase was separated and washed with ethyl acetate (2 x 10 300ml). The aqueous phase was acidified with 5N hydrochloric acid and extracted with ethyl acetate (3x400ml). The organic extracts of the acidic aqueous phase were combined, dried over sodium sulphate, and the solvent evaporated to leave a solid. The solid was dissolved in hot ethyl acetate (100ml) and 40-60 petrol (160ml) was added to 15 the hot solution. The solution was left to cool and the resulting solid filtered off, washed and dried to give the title compound (28.4g)

LC/MS R_t 2.24min m/z 182 [MH^+].

Intermediate 29. 6-({3-[(Dimethylamino)carbonyl]phenyl}thio)-4-[(3-methoxyphenyl)amino]-8-methylquinoline-3-carboxamide




A stirred mixture of Intermediate 45 (0.5g), Intermediate 28 (0.392g), cuprous iodide (0.03g) and potassium carbonate (0.38g) in 1,3-dimethyltetrahydropyrimidin-2(1*H*)-one (7ml) was heated at 100°C for 16h. The mixture was cooled, poured into water (100ml) and extracted into ethyl acetate (3x40ml). The extracts were washed with water (100ml), dried (Na₂SO₄) and evaporated. The residual oil was triturated with ethyl acetate (10ml) to give the title compound as a fawn coloured solid (0.263g).

LC/MS R_t 2.67 min, *m/z* 487 [MH⁺]

10

Intermediate 37. 1,1-Dimethylethyl [(6-iodo-4-{[3-(methoxy)phenyl]amino}-3-quinolinyl)carbonyl]carbamate




15

To a stirred suspension of Intermediate 14 (6.93g) in dichloromethane (170ml) was added *N,N*-dimethyl-4-aminopyridine (2.42g) followed by di-*tert*-butyldicarbonate (18g) (Aldrich). The solution was stirred at room temperature for 25 min, then quenched by addition of aqueous citric acid (200ml) and stirred vigorously for 30 min. The organic layer was separated and the aqueous layer washed with dichloromethane (50ml). The combined organic extracts were washed with brine (100ml), dried over magnesium sulphate and concentrated *in vacuo*. The residue was triturated in diethyl ether to give a yellow solid which was collected by filtration, washed with diethyl ether (3 x 15ml) and dried *in vacuo* to give the title compound as a yellow solid (6.6 g).

LC/MS R_t 3.59min m/z 520 $[MH^+]$

Intermediate 47. 6-[(4-{[*tert*-Butyl(dimethyl)silyl]oxy}phenyl)thio]-4-[(3-methoxyphenyl)amino]quinoline-3-carboxamide




A stirred mixture of Intermediate 37 (0.8g) and 4-[(*tert*-butyl(dimethyl)silyl]oxy} benzenethiol (0.74g, EP465802A1), with (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.05g), potassium *tert*-butoxide (0.26g) and tris(dibenzylideneacetone) dipalladium(0) (0.08g) in toluene (30ml) was heated at 106°C for 18h. The mixture was cooled, diluted with ethyl acetate (25ml) and washed with a sodium carbonate solution (30ml). The organic extracts were washed with water (30ml), dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by chromatography on silica gel using ethyl acetate / diethyl ether (7:3) as the eluent. The appropriate fractions were concentrated *in vacuo* to give the title compound as a yellow foam (0.38g).

LC/MS R_t 3.9min, m/z 532 [MH^+]

Intermediate 51. 1,1-Dimethylethyl 4-amino-1*H*-indazole-1-carboxylate

5



To a solution of 4-nitro-1*H*-indazole (1.57g, *Journal of Heterocyclic Chemistry* 1979, **16(8)**, 1599-603) and di-tert-butyl dicarbonate (2.33g) in acetonitrile (30ml) was added *N,N*-dimethyl-4-aminopyridine (0.059g). The reaction mixture was stirred at room temperature for 30 min, then concentrated *in vacuo* to leave a brown solid which was purified by silica SPE, eluting sequentially with dichloromethane and diethyl ether to give 1,1-dimethylethyl 4-nitro-1*H*-indazole-1-carboxylate as a yellow solid (1.9g).

LC/MS R_t 3.26 min, m/z 263 [MH^+]

15


1,1-Dimethylethyl 4-nitro-1*H*-indazole-1-carboxylate (1.2g) was dissolved in ethanol (150 ml) and stirred with 10% palladium on carbon (0.24g) under an atmosphere of hydrogen (1 atmosphere pressure) for 18h. The solution was filtered through a pad of celite and the filtrate concentrated *in vacuo* to give the title compound as a yellow-orange solid (1.03g).

20

LC/MS R_t 2.36 min, m/z 234 [MH^+]

Intermediate 52: Ethyl 3-[(3-(aminocarbonyl)-4-{[4-fluoro-3-(methoxy)phenyl]amino}-8-methyl-6-quinolinyl)thio]propanoate


25





A mixture containing Intermediate 35 (1.4g), ethyl 3-mercaptopropionate (0.74g, available from Aldrich), potassium *tert*-butoxide (0.64g), tris(dibenzylideneacetone) dipalladium(0) (0.26g) and (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.15g) was dissolved in *N,N*-dimethylformamide (20ml) and stirred under an atmosphere of nitrogen at 100°C for 18h. The solvents were concentrated *in vacuo* and the residue dissolved in methanol. This was purified by chromatography on an SPE column eluting with methanol and a solution of ammonia in methanol, to give the title compound as a brown foam (1.06g).

LC/MS R_t 2.69 min, m/z 458 [MH^+]


Similarly prepared were the following:



Intermediate Number	R ¹ NH-	Starting Material	LCMS MH ⁺	LCMS R _t (min)
Intermediate 57		Intermediate 36	452	2.67

Intermediate 58		Intermediate 62	411	2.36
Intermediate 103		Intermediate 61	428	2.86

Intermediate 53: 3-[(3-(Aminocarbonyl)-4-{[4-fluoro-3-(methoxy)phenyl]amino}-8-methyl-6-quinoliny]thio]propanoic acid




5

A solution of Intermediate 52 (0.95g) in ethanol (10ml) was treated with 2M sodium hydroxide (10ml) and the resulting solution was left standing at room temperature overnight. The solvent was evaporated *in vacuo*. The residue was dissolved in water and
 10 acidified with 2M hydrochloric acid to pH 4. The resulting precipitate was filtered off, washed with water and dried *in vacuo* to give the title compound as an orange solid (0.8g).


LC/MS R_t 2.3 min, m/z 430 [MH^+]

15 Similarly prepared were the following:



Intermediate Number	R ¹ NH-	Starting Material	LCMS MH ⁺	LCMS R _t (min)
Intermediate 59		Intermediate 103	400	2.39
Intermediate 60		Intermediate 58	383	2.10

5 **Intermediate 54.** 4-[[3-(Methoxy)phenyl]amino]-6-(4-piperidinylsulfonyl)-3-quinolinecarboxamide trifluoroacetate




- 10 To a mixture containing Example 377 (0.64g) in anisole (9ml) was added a solution of 95% trifluoroacetic acid in water (16ml). The mixture was stirred for 1.5h at room temperature and was then concentrated *in vacuo*. The residue was co-evaporated with

toluene (2 x 20ml), triturated with ethyl acetate and filtered to give a yellow solid. This residue was again triturated with ethyl acetate and filtered to give the title compound as a yellow solid (0.570g).

LC/MS R_t 1.94 min, m/z 455 $[MH^+]$

5

Intermediate 55. 1,1-Dimethylethyl 4-[(3-(aminocarbonyl)-4-{[4-fluoro-3-(methoxy)phenyl]amino}-8-methyl-6-quinoliny]sulfonyl]-1-piperidinecarboxylate



10

To a solution of Intermediate 35 (1.0g) in *N,N*-dimethylformamide (30ml) under an atmosphere of nitrogen was added 1,1-dimethylethyl 4-mercapto-1-piperidinecarboxylate (0.89g, US5317025A), potassium *tert*-butoxide (0.46g), tris(dibenzylideneacetone) dipalladium(0) (0.19g) and (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.11g). The mixture was heated to 100°C for 3h, cooled and the solvent removed under reduced pressure. The residue was partitioned between ethyl acetate (100ml) and water (100ml) then dried over magnesium sulphate, filtered and concentrated *in vacuo*. The residue was purified by SPE (eluting with a gradient of 0 to 5% methanol in chloroform) to give intermediate 1,1-dimethylethyl 4-[(3-(aminocarbonyl)-4-{[4-fluoro-3-(methoxy)phenyl]amino}-8-methyl-6-quinoliny]thio]-1-piperidinecarboxylate as a yellow solid (1.1g). This sulphide was dissolved in *N,N*-dimethylformamide (50ml) and oxone (5.15g) was added portionwise. The mixture was stirred at room temperature for 3h, then quenched by addition of 1M sodium sulphite solution (500ml). The mixture was extracted with chloroform (2 x 200ml), and the organic layers were washed with 10% lithium chloride solution, dried over magnesium sulphate, filtered and concentrated to give the title compound as a pale yellow solid (0.71g) after trituration with ether.


25

LC/MS R_t 3.04 min, m/z 573 $[MH^+]$

Similarly prepared from Intermediate 35 and 1,1-dimethylethyl (2-mercaptoethyl) carbamate (available from Aldrich) was :

5


Intermediate 56. 1,1-Dimethylethyl {2-[(3-(aminocarbonyl)-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-6-quinolinyl)sulfonyl]ethyl}carbamate



10

LC/MS R_t 2.79 min, m/z 533 $[MH^+]$

Intermediate 62. 6-Iodo-8-methyl-4-(3-pyridinylamino)-3-quinolinecarboxamide hydrochloride




15

To a solution of Intermediate 48 (1.1g) in *N,N*-dimethylformamide (20ml) was added 3-aminopyridine (0.8g, available from Aldrich) and pyridine hydrochloride (0.7g, available from Aldrich). The mixture was heated at 80°C under nitrogen for 2 days. The solvent was evaporated *in vacuo*. The residue was triturated with methanol and the precipitate filtered off to give the title compound as a brown solid (0.9g).

20








LC/MS R_t 2.32min m/z 405 $[MH^+]$.

Similarly prepared were the following:



5

Intermediate Number (a) (b)	R ¹ NH-	R ²⁰ -	Starting Material	Amine / Supplier	LCMS MH ⁺	LCMS R _t (min)
Intermediate 81 HCl		Et-	Intermediate 72	3-pyridinamine / Aldrich	419	2.52
Intermediate 82 HCl		Et-	Intermediate 72	5-chloro-3-pyridinamine / Synchem OHG	453	3.19
Intermediate 83 HCl		Et-	Intermediate 72	5-fluoro-3-pyridinamine / Synchem OHG	437	2.93
Intermediate 84 HCl		F-	Intermediate 87	5-fluoro-3-pyridinamine / Synchem OHG	427	2.6
Intermediate 85 HCl		F-	Intermediate 87	5-chloro-3-pyridinamine / Synchem OHG	443	2.88
Intermediate 86 HCl		F-	Intermediate 87	3-pyridinamine / Aldrich	409	2.27


96 HCl		Cl-	Intermediate 49	3-pyridinamine / Aldrich	425	2.52
97 HCl		Me-	Intermediate 48	5-chloro-3- pyridinamine / Synchem OHG	439	2.95
98 HCl		Me-	Intermediate 48	5-fluoro-3- pyridinamine / Synchem OHG	423	2.65
99 HCl		Cl-	Intermediate 49	5-fluoro-3- pyridinamine / Synchem OHG	443	2.91
100 HCl		Cl-	Intermediate 49	5-chloro-3- pyridinamine / Synchem OHG	459	2.94
101 HCl		Me-	Intermediate 48	1-ethyl-1H- pyrazol-5-amine / Aldrich	422	2.58
102 HCl		Cl-	Intermediate 49	1-ethyl-1H- pyrazol-5-amine / Aldrich	442	2.86

(a) Salt form HCl = hydrochloride

(b) All products isolated by trituration with acetonitrile and filtration.

5


Intermediate 73. 2-Ethyl-4-iodoaniline



To a stirred solution of 2-ethylaniline (1.88g, available from Aldrich) and sodium acetate (1.27g) in acetic acid (20ml) was added iodine monochloride (1ml, available from Aldrich). The mixture was stirred at 20°C for 90 min and then the solvent was removed *in vacuo*. The residue was partitioned between ethyl acetate (25ml) and saturated aqueous sodium carbonate solution (25ml). The organic layer was dried using a hydrophobic frit and the solvent was removed *in vacuo*. Purification by C18 SPE eluting with 20% acetonitrile in water gave the title compound as a purple solid (0.402g).

LC/MS R_t 3.23 min, m/z 248 $[MH^+]$


Intermediate 64. 7-({3-[(Dimethylamino)carbonyl]phenyl}thio)-6-iodo-4-({3-(methoxy)phenyl}amino)-3-quinolinecarboxamide



A stirred mixture of Intermediate 63 (0.4g), Intermediate 28 (0.16g) and potassium carbonate (0.38g) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (10ml) was heated at 100°C under nitrogen for 3h. A further portion of Intermediate 28 (0.07g) was added and the mixture stirred at 60°C for 23h. The cooled mixture was diluted with water (100ml) and extracted with ethyl acetate (3 x 100ml). The combined organic extracts were washed with water (2 x 70ml) and brine (70ml), dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified by chromatography on silica gel eluting with methanol followed by mass directed preparative HPLC (Method A) to give the title compound as a yellow foam (0.12g).

LC/MS R_t 2.94min, m/z 599 $[MH^+]$

Intermediate 65. 7-({3-[(Dimethylamino)carbonyl]phenyl}sulfinyl)-6-iodo-4-{3-(methoxy)phenylamino}-3-quinolinecarboxamide



5


Oxone (0.5g) was added portionwise to a stirred solution of Intermediate 64 (0.12g) in *N,N*-dimethylformamide (5ml). The solution was stirred at room temperature under nitrogen for 21h. More oxone (0.5g) was added and the mixture was stirred for a further 3h, quenched with a solution of sodium sulphite (1.5g) in water (15ml), diluted with water (50ml) and extracted with ethyl acetate (3 x 50ml). The combined organic extracts were dried over magnesium sulphate and concentrated *in vacuo* to give the title compound as a yellow solid (0.15g).

10

LC/MS R_t 2.70min, m/z 615 $[MH^+]$

15

Intermediate 69. 5-Mercapto-*N,N*-dimethyl-3-pyridinecarboxamide



Sodium thiomethoxide (3g) was added to a stirred solution of 5-bromo-*N,N*-dimethyl-3-pyridinecarboxamide (2.5g, WO2000055168) in *N,N*-dimethylformamide (40ml) and the suspension stirred at 100°C for 4h. The solvent was concentrated *in vacuo*, the residue dissolved in 2M sodium hydroxide (35ml) and water (50ml), and the solution washed with chloroform (4 x 75ml). The aqueous layer was acidified with 2M hydrochloric acid to pH 4 and extracted with chloroform (5 x 80ml), and the combined organic layers were washed


25

with brine (20ml), dried over magnesium sulphate and concentrated *in vacuo* to give the title compound as an orange oil (1.8g).

LC/MS R_t 0.96min, m/z 183 $[MH^+]$

5

Intermediate 70. 1-Methyl-4-nitro-2,3-dihydro-1H-indole




To a stirred solution of 1-methyl-4-nitro-1H-indole (3.8g, Organic Process Research and Development 2001 5 (6) 604) and borane-tetrahydrofuran complex (1M in tetrahydrofuran, 86.3ml) at 0°C under an atmosphere of nitrogen was added, dropwise, trifluoroacetic acid (88ml). The resulting mixture was allowed to warm to room temperature and stirred at room temperature for 90 min. The mixture was cautiously added to 2M sodium carbonate solution (750ml) over 20 min and then stirred for 30 min. The mixture was extracted with ethyl acetate (2x300ml), the combined organic extracts were dried over sodium sulphate and the solvent was removed *in vacuo*. Purification by column chromatography on silica gel, eluting with hexane : ethyl acetate (9:1) gave the title compound as a red solid (1.97g).

TLC SiO_2 (hexane : ethyl acetate (4:1)) R_f = 0.61

20

Intermediate 71. 1-Methyl-2,3-dihydro-1H-indol-4-amine




A solution of Intermediate 70 (0.50g) in ethanol (30ml) was added to 10% palladium on carbon (0.050g) and the mixture was stirred under an atmosphere of hydrogen for 20 min. The mixture was filtered through 'hyflo' filter aid and the solvent removed *in vacuo* to give the title compound as a brown oil (0.405g).

5 TLC SiO₂ (hexane : ethyl acetate (4:1)) R_f = 0.25

Examples


Experimental details for the preparation of representative Examples are given in full below. Summary details for further Examples prepared by analogous methods are given in the accompanying tables.

5










Example 10. 4-[(3-Fluorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide








- 10 Intermediate 9 (0.014g) was suspended in acetonitrile (3ml), 3-fluoroaniline (0.0056g, available from Aldrich) was added, and the mixture was heated under reflux for 16h. After cooling to room temperature, the mixture was cooled in a refrigerator for 2h, filtered, and the residue purified by mass directed preparative HPLC (Method A) to give the title compound (0.011g).
- 15 LC/MS R_t 1.95min m/z 359 [MH^+]







Similarly prepared were the following:












Ex. No.	R^1NH-	R^3SO_2-	Starting Material	Amine Reagent/Source	Isolation Method	LCMS MH^+	LCMS R_t
(a)					(b)		(min)








1 HCl		MeSO ₂ ⁻	Intermediate 9	6-aminobenzothiazole/ Lancaster	(I)	399	1.9
2 HCl		MeSO ₂ ⁻	Intermediate 9	3-amino-N-methylacetanilide/ Merlin synthesis	(I)	413	1.89
3 HCl		MeSO ₂ ⁻	Intermediate 9	N,N-dimethylbenzene-1,3-diamine hydrochloride / Aldrich	(I)	385	1.83
4 HCl		MeSO ₂ ⁻	Intermediate 9	3,5-dimethoxyaniline/ Aldrich	(I)	402	2.05
5 HCl		MeSO ₂ ⁻	Intermediate 9	1,2-benzisoxazol-5-amine/ Key organics Ltd (8W-0024)	(I)	383	1.84
6 HCl		MeSO ₂ ⁻	Intermediate 9	methyl 3-amino-4-oxobenzoate hydrochloride/ Fluka	(I)	400	1.99
7 HCl		MeSO ₂ ⁻	Intermediate 9	3-methylaniline hydrochloride/ TCI-JP	(I)	356	2
8 HCl		MeSO ₂ ⁻	Intermediate 9	3-aminobenzonitrile/ Aldrich	(I)	367	2.26
9		MeSO ₂ ⁻	Intermediate 9	3-aminobenzyl alcohol/ Aldrich	(II)	372	1.82










11		MeSO ₂ -	Intermediate 9	1-methyl-1H-benzimidazol-6-amine/ <i>Heterocycles</i> , 1991, 32 (5), 1003-12.	(II)	396	1.65
12		MeSO ₂ -	Intermediate 9	3-chloro-4-fluoroaniline/ ABCR	(II)	394	2.37
13 HCl		MeSO ₂ -	Intermediate 9	aniline/ Aldrich	(I)	342	2.07
15 HCl		MeSO ₂ -	Intermediate 9	6-aminobenzoxazinone/ WO 9845268 A1	(I)	399	1.99
16 HCl		MeSO ₂ -	Intermediate 9	2,3-dihydro-1H-inden-5-ylamine hydrochloride/ Aldrich	(I)	382	2.48
17 HCl		MeSO ₂ -	Intermediate 9	2,3-dihydro-1,4-benzodioxin-6-amine hydrochloride/ Aldrich	(I)	400	2.18
18 HCl		MeSO ₂ -	Intermediate 9	2,3-dihydro-1,4-benzodioxin-5-amine hydrochloride/ WO 9703067 A1	(I)	400	2.2








19 HCl		MeSO ₂ ⁻	Intermediate 9	1-amino-5,6,7,8-tetrahydronaphthalene/ Aldrich	(I)	396	2.57
20 HCl		MeSO ₂ ⁻	Intermediate 9	2,3-dihydro-1-benzofuran-4-amine hydrobromide/ <i>Journal of Heterocyclic Chemistry</i> , 1980, 17 (6), 1333-5.	(I)	384	2.23
21		MeSO ₂ ⁻	Intermediate 9	7-amino-1,3-benzoxazol-2(3H)-one/ <i>Annales Universitatis Mariae Curie-Skłodowska, Sectio D: Medicina</i> , 1980, Volume Date 1979, 35 121-8.	(II)	399	1.93
22 HCl		MeSO ₂ ⁻	Intermediate 9	3-ethylaniline/ Aldrich	(I)	370	2.33
23 HCl		MeSO ₂ ⁻	Intermediate 9	3-isopropylaniline/ APIN	(I)	384	2.5
24 HCl		MeSO ₂ ⁻	Intermediate 9	3-chloro-4-methoxyaniline / Aldrich	(I)	406/408	2.2








25		MeSO ₂ ⁻	Intermediate 9	3-(methoxymethyl)aniline/ WO 0018721 A1	(II)	386	2.01
26 HCl		MeSO ₂ ⁻	Intermediate 9	2,5-dimethoxyaniline/ Aldrich	(I)	402	2.37
27 HCl		MeSO ₂ ⁻	Intermediate 9	3-methoxyaniline / Aldrich	(I)	372	2.33
28 HCl		MeSO ₂ ⁻	Intermediate 9	3-hydroxy-4-methoxyaniline / Aldrich	(I)	388	2.0
29 HCl		MeSO ₂ ⁻	Intermediate 9	3-phenoxyaniline / Aldrich	(I)	434	2.97
30 HCl		MeSO ₂ ⁻	Intermediate 9	3-amino-2-methylphenol/ Aldrich	(I)	372	2.2
31 HCl		MeSO ₂ ⁻	Intermediate 9	3-aminophenol hydrochloride/ TCI-US	(I)	358	2.04
32 HCl		MeSO ₂ ⁻	Intermediate 9	4-fluoro-3-methoxyaniline / Apollo-Chem	(I)	390	1.92
33 HCl		MeSO ₂ ⁻	Intermediate 9	2-(3-amino phenoxy) ethanol hydrochloride/ J. Amer. Chem. Soc.; 1937, 59; 1716	(I)	402	1.8

34 HCl		MeSO ₂ ⁻	Intermediate 9	3-ethoxyaniline/ Aldrich	(I)	386	2.04
35 HCl		MeSO ₂ ⁻	Intermediate 9	3-chloroaniline/ Aldrich	(I)	376	2.09
36 HCl		MeSO ₂ ⁻	Intermediate 9	[3-(2-methoxyethoxy)phenyl] amine hydrochloride/ EP 388165 A2	(I)	416	1.94
37 HCl		MeSO ₂ ⁻	Intermediate 9	4-methoxyaniline / Aldrich	(I)	372	1.88
38 HCl		MeSO ₂ ⁻	Intermediate 9	3,4-dimethoxy aniline/ Aldrich	(I)	402	1.84
39		PhSO ₂ ⁻	Intermediate 8	3-ethoxyaniline/ Aldrich	(II)	448	2.55
40		PhSO ₂ ⁻	Intermediate 8	[3-(2-methoxyethoxy)phenyl] amine hydrochloride/ EP 388165 A2	(II)	478	1.42
41		PhSO ₂ ⁻	Intermediate 8	3-amino-N-methyl benzamide/ TCI-US	(II)	461	2.37












42		PhSO ₂ -	Intermediate 8	4-aminophenol hydrochloride/ Aldrich	(II)	420	2.46
43 HCl		PhSO ₂ -	Intermediate 8	6-aminobenzothiazole/ Lancaster	(I)	461	2.67
44 HCl		PhSO ₂ -	Intermediate 8	N-(3-aminophenyl)- N-methyl acetamide/ Merlin Synthesis	(I)	475	2.58
45 HCl		PhSO ₂ -	Intermediate 8	1-methyl-1H-benzimidazol- 6-amine/ Gwyn Ellis <i>Heterocycles</i> , 1991, 32(5) , 1003- 12.	(I)	458	2.24
46 HCl		PhSO ₂ -	Intermediate 8	3,5-dimethoxy aniline/ Aldrich	(I)	464	2.89
47 HCl		PhSO ₂ -	Intermediate 8	methyl 3-amino benzoate hydrochloride/ Fluka	(I)	462	2.86
48 HCl		PhSO ₂ -	Intermediate 8	4-(2-morpholin-4-ylethoxy) aniline EP 410358 A1	(I)	533	2.1



















49 HCl		PhSO ₂ -	Intermediate 8	3-fluoroaniline/ Aldrich	(I)	422	2.96
50 HCl		PhSO ₂ -	Intermediate 8	3-chloro-4-fluoroaniline/ Aldrich	(I)	456	3.12
51 HCl		PhSO ₂ -	Intermediate 8	3-methylaniline hydrochloride/ TCI- US	(I)	418	2.87
52 HCl		PhSO ₂ -	Intermediate 8	3-amino benzonitrile/ Aldrich	(I)	429	2.91
53		PhSO ₂ -	Intermediate 8	N-(4-aminophenyl) morpholine-4-carboxamide / Peakdale Molecular Ltd	(II)	532	2.34
54		PhSO ₂ -	Intermediate 8	N,N-dimethyl benzene-1,3-diamine hydrochloride / Aldrich	(II)	447	2.75
55		PhSO ₂ -	Intermediate 8	N-(3-aminophenyl) acetamide hydrochloride / Acros Chimica	(II)	461	2.4
56		PhSO ₂ -	Intermediate 8	3-aminobenzyl alcohol/ Aldrich	(II)	434	2.34
57 HCl		PhSO ₂ -	Intermediate 8	aniline/ Aldrich	(I)	404	2.73

58 HCl		PhSO ₂ -	Intermediate 8	1-acetylindoline-5-amine / Maybridge	(I)	487	2.52
59 HCl		PhSO ₂ -	Intermediate 8	6-amino-1,3-benzoxazol-2(3H)-one / WO 9845268 A1	(I)	461	2.48
60 HCl		PhSO ₂ -	Intermediate 8	1-acetyl-6-aminoindoline / SIGMA	(I)	487	2.55
61 HCl		PhSO ₂ -	Intermediate 8	6-amino-2,3-dihydro-1H-inden-1-one / J. Med. Chem. 2003, 46(3), 399-408.	(I)	458	2.67
62 HCl		PhSO ₂ -	Intermediate 8	6-amino-1,2,3,4-tetrahydronaphthalen-1-one / Maybridge	(I)	472	2.93
63 HCl		PhSO ₂ -	Intermediate 8	2,3-dihydro-1,4-benzodioxin-6-amine hydrochloride / Aldrich	(I)	462	2.69
64 HCl		PhSO ₂ -	Intermediate 8	1-amino-5,6,7,8-tetrahydronaphthalene / Aldrich	(I)	458	3.07








65 HCl		PhSO ₂ -	Intermediate 8	7-amino-1,3-benzoxazol-2(3H)-one/ <i>Medicina</i> 1980, Volume Date 1979, 35 , 121-8.	(I)	461	2.64
66 HCl		PhSO ₂ -	Intermediate 8	2,3-dihydro-1-benzofuran-4-amine hydrobromide/ <i>Journal of Heterocyclic Chemistry</i> 1980, 17(6) , 1333-5.	(I)	446	2.84
67		PhSO ₂ -	Intermediate 8	2,3-dihydro-1H-inden-5-amine hydrochloride/ Aldrich	(II)	444	2.95
68		PhSO ₂ -	Intermediate 8	2,3-dihydro-1,4-benzodioxin-5-amine hydrochloride/ WO 9703067 A1	(II)	462	2.69
69 HCl		PhSO ₂ -	Intermediate 8	3-ethylaniline/ Aldrich	(I)	432	2.92
70 HCl		PhSO ₂ -	Intermediate 8	3-isopropylaniline/ APIN	(I)	446	3.04
71 HCl		PhSO ₂ -	Intermediate 8	3-chloro-4-methoxyaniline / Aldrich	(I)	468	2.76







72 HCl		PhSO ₂ -	Intermediate 8	8-amino-3,4-dihydro-1(2H)-naphthalenone / WO 0160826 A2	(I)	472	2.86
73		PhSO ₂ -	Intermediate 8	3-[(methyloxy)methyl]aniline / WO 0018721 A1	(II)	448	2.60
74 HCl		PhSO ₂ -	Intermediate 8	3-(methyloxy)aniline / Aldrich	(I)	434	2.77
75 HCl		PhSO ₂ -	Intermediate 8	2-[(3-aminophenyl)oxy]ethanol hydrochloride / J. Amer. Chem. Soc.; 1937, 59 , 1716	(I)	464	2.21
76 HCl		PhSO ₂ -	Intermediate 8	4-amino-N-methylbenzamide / Buttpark	(I)	461	2.2
77		PhSO ₂ -	Intermediate 8	3-amino-2-methylphenol / Aldrich	(II)	434	2.57
78 HCl		PhSO ₂ -	Intermediate 8	4-methoxyaniline hydrochloride / Acros	(I)	434	2.32
79 HCl		PhSO ₂ -	Intermediate 8	3-[(trifluoromethyl)oxy]aniline / Aldrich	(I)	487	2.8








80		PhSO ₂ -	Intermediate 8	3-(4-morpholinyl)aniline/ <i>Journal of Organic Chemistry</i> 2002, 67 (9), 3029-3036.	(II)	489	2.72
81		PhSO ₂ -	Intermediate 8	3-aminophenol hydrochloride/ TCI-US	(II)	420	2.6
82		PhSO ₂ -	Intermediate 8	[3,4-bis(methoxy)phenyl]amine hydrochloride/ Aldrich	(II)	463	2.58
83 HCl			Intermediate 7	6-aminobenzo thiazole/ Lancaster	(I)	453	2.3
84 HCl			Intermediate 7	N-(3-aminophenyl)- N-methyl acetamide/ Merlin synthesis	(I)	467	2.29
85 HCl			Intermediate 7	N-(4-aminophenyl)- 4-morpholine carboxamide/ Peakdale molecular Ltd	(I)	524	2.21
86 HCl			Intermediate 7	3,5-dimethoxy aniline/ Aldrich	(I)	456	2.51








87 HCl			Intermediate 7	methyl 3-amino benzoate hydrochloride/ Fluka	(I)	454	2.48
88 HCl			Intermediate 7	3-amino-N-methyl benzamide/ TCI-US	(I)	453	2.17
89 HCl			Intermediate 7	3-fluoroaniline/ Aldrich	(I)	414	2.53
90 HCl			Intermediate 7	3-chloro-4-fluoroaniline/ Aldrich	(I)	448	2.72
91 HCl			Intermediate 7	(3-methylphenyl) amine hydrochloride/ TCI-US	(I)	410	2.49
92			Intermediate 7	N,N-dimethyl-1,3-benzene diamine hydrochloride/ Aldrich	(II)	439	2.62
93			Intermediate 7	N-(3-aminophenyl) acetamide hydrochloride/ Acros	(II)	453	2.25
94			Intermediate 7	2-chloroaniline/ Aldrich	(II)	430	2.69
95 HCl			Intermediate 7	aniline/ Aldrich	(I)	396	2.58








96 HCl			Intermediate 7	1-methyl-1H-benzimidazol-6-amine/ <i>Heterocycles</i> 1991, 32(5) , 1003-12.	(I)	450	2.04
97 HCl			Intermediate 7	3-ethylaniline/ Aldrich	(I)	424	2.81
98 HCl			Intermediate 7	3-isopropylaniline/ TCI-US	(I)	438	3.0
99 HCl			Intermediate 7	3-methoxyaniline/ Aldrich	(I)	426	2.58
100		PhSO ₂ -	Intermediate 8	3-aminopyridine/ Aldrich	(II)	405	2.41
145 HCl		MeSO ₂ -	Intermediate 9	4-fluoro-3-methylaniline/ Aldrich	(I)	396	1.77
146 TFA		MeSO ₂ -	Intermediate 9	5-amino-2-methylbenzofuran hydrochloride/ Aldrich	(III)	396	2.43
147 TFA		MeSO ₂ -	Intermediate 9	1-methyl-1H-indazol-6-amine hydrochloride/ <i>Synthetic Communications</i> , 1996, 26(13) , 2443-2447.	(III)	396	2.15







148 TFA		MeSO ₂ ⁻	Intermediate 9	1-methyl-1 <i>H</i> -indazole-5-amine/ Bionet Research Ltd	(III)	396	2.06
149 TFA		MeSO ₂ ⁻	Intermediate 9	1-(phenylmethyl)-1 <i>H</i> -indazole-5-amine/ WO 0283654 A1	(III)	472	2.56
150 HCl		MeSO ₂ ⁻	Intermediate 9	3-(trifluoromethyl)aniline/ Aldrich	(I)	410	2.05
151 HCl		MeSO ₂ ⁻	Intermediate 9	1-(3-aminophenyl)ethanone/ Aldrich	(I)	384	1.71
152 HCl		MeSO ₂ ⁻	Intermediate 9	3-(5-methyl-1,2,4-oxadiazol-3-yl)aniline/ Intermediate 19	(I)	424	1.88
153 HCl		MeSO ₂ ⁻	Intermediate 9	1-(3-aminophenyl)- <i>N,N</i> -dimethylmethanesulfonamide/ Peakdale molecular Ltd	(I)	463	1.78
154 HCl		MeSO ₂ ⁻	Intermediate 9	3-(3-thienyl)aniline/ US 6211220 B1	(I)	424	2.15

155 HCl		MeSO ₂ ⁻	Intermediate 9	3-methyl-2,3-dihydro-1-benzofuran-4-amine/ Intermediate 25	(I)	398	1.91
156		MeSO ₂ ⁻	Intermediate 9	4-amino-2-methyl-1H-isoindole-1,3(2H)-dione/ <i>Archiv der Pharmazie</i> (Weinheim, Germany) 1989, 322(7) , 419-26.	(II)	425	2.35
157 TFA		PhSO ₂ ⁻	Intermediate 8	1,2-benzisoxazol-5-amine / Key organics Ltd	(III)	445	2.61
158 TFA		PhSO ₂ ⁻	Intermediate 8	1,2-dimethyl-1H-benzimidazol-6-amine/ Intermediate 27	(III)	472	2.15
159 TFA		PhSO ₂ ⁻	Intermediate 8	3,5-dichloroaniline/ Aldrich	(III)	472	3.35
160 TFA		PhSO ₂ ⁻	Intermediate 8	2-methyl-1,3-benzoxazol-5-amine/ <i>Collection of Czechoslovak Chemical Communications</i> 1996, 61(3) , 371-	(III)	459	2.33

				380.			
161 TFA		PhSO ₂ -	Intermediate 8	4-chloro-3-methoxyaniline / Wychem	(III)	468	3.08
162 TFA		PhSO ₂ -	Intermediate 8	5-amino-2-methylbenzofuran hydrochloride/ Sigma Aldrich	(III)	458	2.92
163 TFA		PhSO ₂ -	Intermediate 8	1-methyl-1H-indazol-6-amine hydrochloride/ Heterocycles, 1995, 41(3), 487-96.	(III)	458	2.67
164 TFA		PhSO ₂ -	Intermediate 8	5-fluoro-2-methoxyaniline / Wychem	(III)	452	2.92
165 TFA		PhSO ₂ -	Intermediate 8	1-methyl-1H-indazol-5-amine/ Bionet Research Ltd	(III)	458	2.51
166 TFA		PhSO ₂ -	Intermediate 8	1-(phenylmethyl)-1H-indazol-5-amine/ WO 0283654 A1	(III)	534	2.97
167 TFA		PhSO ₂ -	Intermediate 8	4-amino-2-(methoxy)phenol hydrochloride/ Journal of Chemical	(III)	450	2.42

				Research, Synopsis 1988, (9), 284-5.			
168 TFA		PhSO ₂ -	Intermedia te 8	3-fluoro-4- methylaniline/ Aldrich	(III)	436	3.04
169 TFA		PhSO ₂ -	Intermedia te 8	2-methyl-1,3- benzothiazol- 6-amine/ AsInEx Compound Collection	(III)	475	2.73
170 TFA		PhSO ₂ -	Intermedia te 8	1H-indol-6- amine/ Lancaster Synthesis	(III)	443	2.7
171 TFA		PhSO ₂ -	Intermedia te 8	3-chloro-5- (methoxy) aniline/ J. Chem. Soc. Perkin 2, 1977, 14.	(III)	468	3.17
172 HCl		PhSO ₂ -	Intermedia te 8	3-(2-methyl-4- pyrimidinyl) aniline/ Fluorochem	(I)	496	2.23
173 HCl		PhSO ₂ -	Intermedia te 8	3-(trifluoro methyl)aniline / Aldrich	(I)	472	2.62
174 HCl		PhSO ₂ -	Intermedia te 8	1-(3- aminophenyl) ethanone/ Aldrich	(I)	446	2.2

175 HCl		PhSO ₂ -	Intermediate 8	3-(1,3-oxazol-5-yl)aniline/ Fluorochem	(I)	471	2.24
176 HCl		PhSO ₂ -	Intermediate 8	3-(1-methyl-1H-pyrazol-3-yl)aniline/ Intermediate 26	(I)	484	2.27
177 HCl		PhSO ₂ -	Intermediate 8	3-(5-methyl-1,2,4-oxadiazol-3-yl)aniline/ Intermediate 19	(I)	486	2.34
178 HCl		PhSO ₂ -	Intermediate 8	1-(3-aminophenyl)- N,N-dimethyl methane sulfonamide/ Peakdale molecular Ltd	(I)	525	2.23
179 HCl		PhSO ₂ -	Intermediate 8	3-(3-thienyl)aniline/ US 6211220 B1	(I)	486	2.55
180 HCl		PhSO ₂ -	Intermediate 8	3-methyl-2,3-dihydro-1-benzofuran-4-amine/ Intermediate 25	(I)	460	2.36
181 HCl		PhSO ₂ -	Intermediate 8	2,2-dimethyl-2,3-dihydro-1-benzofuran-7-amine/ DE 3526510 A1	(I)	474	2.34


182		PhSO ₂ ⁻	Intermediate 8	4-amino-2-methyl-1H-indole-1,3(2H)-dione/ <i>Archiv der Pharmazie</i> (Weinheim, Germany) 1989, 322(7) , 419-26.	(II)	487	2.85
183		PhSO ₂ ⁻	Intermediate 8	[4-(methoxy)-2-naphthalenyl]amine 4-methylbenzene sulfonate/ Sigma	(I)	484	3.19
590 HCl			Intermediate 95	5-chloro-3-pyridinamine / Synchem OHG	(I)	419	2.44
589 HCl			Intermediate 95	3-pyridinamine / Aldrich	(I)	385	1.96

(a) Salt forms: HCl = hydrochloride
TFA = trifluoroacetate












(b) Isolation Method:

- 5 (I) Filtered off directly from the reaction mixture.
(II) Mass Directed preparative HPLC Method A.
(III) Mass Directed preparative HPLC Method B.

The following compounds were prepared by a similar method to Example 10:

















Ex. No. (a)	R ¹ NH-	R ³ SO ₂ -	R ²⁰ -	Starting Material	Amine Reagent/ Source	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
187 HCl		MeSO ₂ -	H-	Intermediate 9	1,1-dimethylethyl (3-aminophenyl) carbamate / <i>J. Med. Chem.</i> 2003, 46 (9) 1661-1669	(I)	457	2.47
188 HCl		MeSO ₂ -	H-	Intermediate 9	1,1-dimethylethyl [(3-aminophenyl)methyl]carbamate / <i>J. Med. Chem.</i> 2003, 46 (9) 1661-1669	(I)	471	2.39
189		MeSO ₂ -	H-	Intermediate 9	1,3-benzodioxol-5-amine / Aldrich	(II)	386	1.98
190 HCl		MeSO ₂ -	H-	Intermediate 9	3-(1,3-oxazol-5-yl)aniline / Maybridge	(I)	409	1.82
191 HCl		MeSO ₂ -	H-	Intermediate 9	3-(3-aminophenyl)-1-methyl-1H-pyrazole / Buttpark	(I)	422	1.88
192 HCl		MeSO ₂ -	H-	Intermediate 9	3-chloro-2-fluoroaniline / Aldrich	(I)	394	2.46
193 HCl		MeSO ₂ -	H-	Intermediate 9	2,3-difluoroaniline / Aldrich	(I)	378	2.3








194 HCl		MeSO ₂ ⁻	H-	Intermediate 9	3-aminobenzonitrile / Aldrich	(I)	367	2.07
195 HCl		MeSO ₂ ⁻	H-	Intermediate 9	5-amino-2-fluorobenzonitrile / Maybridge	(I)	385	2.15
196 HCl		MeSO ₂ ⁻	H-	Intermediate 9	3-isopropoxyaniline / Maybridge	(I)	400	2.35
197 HCl		MeSO ₂ ⁻	H-	Intermediate 9	2-amino-5-fluorotoluene / Aldrich	(I)	374	2.2
198 HCl		MeSO ₂ ⁻	H-	Intermediate 9	3,4-difluoroaniline / Aldrich	(I)	378	2.26
199 HCl		MeSO ₂ ⁻	H-	Intermediate 9	4-fluoro-3-(trifluoromethyl) aniline / Avocado	(I)	428	2.61
200 HCl		MeSO ₂ ⁻	H-	Intermediate 9	2-fluoroaniline / Aldrich	(I)	360	2.04
201 HCl		MeSO ₂ ⁻	H-	Intermediate 9	2,4-difluoroaniline / Aldrich	(I)	378	2.22
202 HCl		MeSO ₂ ⁻	H-	Intermediate 9	2-chloro-4-fluoroaniline / Aldrich	(I)	394	2.43
203 HCl		MeSO ₂ ⁻	H-	Intermediate 9	3-aminopyridine / Aldrich	(IV)	343	1.69
204 HCl		MeSO ₂ ⁻	H-	Intermediate 9	4-aminobenzoic acid / Aldrich	(I)	386	2










205 TFA		MeSO ₂ ⁻	H-	Intermediate 9	4-chloro-3-methoxyaniline / Wychem	(III)	406	2.49
206 TFA		MeSO ₂ ⁻	H-	Intermediate 9	1-methyl-1H-benzimidazol-6-amine/ <i>Heterocycles</i> . 1991, 32 (5), 1003-12.	(III)	396	1.77
207 TFA		MeSO ₂ ⁻	H-	Intermediate 9	6-(methoxy)-1,3-benzothiazol-4-amine <i>J. Am. Chem. Soc.</i> , 1939, 61 (8), 2013-2017.	(III)	429	2.35
208 TFA		MeSO ₂ ⁻	H-	Intermediate 9	3-fluoro-5-(3-pyridinyl)aniline <i>J. Med. Chem.</i> 2000, 43 (6), 1123-1134.	(III)	437	2.34
209 TFA		MeSO ₂ ⁻	H-	Intermediate 9	5-fluoro-2-methoxyaniline / Wychem	(III)	390	2.3
210 TFA		MeSO ₂ ⁻	H-	Intermediate 9	1-methyl-1H-1,2,3-benzotriazol-5-amine/ US 2003060453 A1	(III)	397	1.98
211 TFA		MeSO ₂ ⁻	H-	Intermediate 9	3,5-difluoroaniline / Aldrich	(III)	378	2.51
212 TFA		MeSO ₂ ⁻	H-	Intermediate 9	3-fluoro-4-methylaniline / Aldrich	(III)	374	2.43
213 TFA		MeSO ₂ ⁻	H-	Intermediate 9	4-amino-2-fluorophenol / Apollo	(III)	376	2.01
















214 TFA		MeSO ₂ ⁻	H-	Intermediate 9	4-(methoxy)-2-naphthalenamine/ Sigma	(III)	422	2.73
215 TFA		MeSO ₂ ⁻	H-	Intermediate 9	6-aminoindole / Lancaster	(III)	381	2.25
216 TFA		MeSO ₂ ⁻	H-	Intermediate 9	methyl 4-amino-2-methoxybenzoate / Avocado	(III)	430	2.36
217 HCl		MeSO ₂ ⁻	H-	Intermediate 9	1,3-benzodioxol-4-amine <i>J. Med. Chem.</i> 1979, 22 (11), 1354-7.	(I)	386	2.06
218 HCl		PhSO ₂ ⁻	H-	Intermediate 8	3-aminobenzonitrile / Aldrich	(I)	429	2.80
219		PhSO ₂ ⁻	H-	Intermediate 8	1,3-benzodioxol-5-amine / Aldrich	(II)	448	2.63
220		PhSO ₂ ⁻	H-	Intermediate 8	3-amino-N,N-dimethylbenzene sulfonamide/ WO 9737646 A1	(II)	511	2.94
221		PhSO ₂ ⁻	H-	Intermediate 8	3-aminobenzene sulfonamide / Fluka	(II)	483	2.6
222		PhSO ₂ ⁻	H-	Intermediate 8	4-amino-N-methylbenzene-sulfonamide / Zelinsky BB	(II)	497	2.51

223 TFA		PhSO ₂ -	H-	Intermediate 8	6-(methoxy)- 1,3-benzothiazol- 4-amine <i>J. Am. Chem. Soc.</i> , 1939, 61(8) , 2013- 2017.	(III)	491	2.92
224 TFA		PhSO ₂ -	H-	Intermediate 8	3-fluoro-5-(3- pyridinyl)aniline / <i>J. Med. Chem.</i> , 2000, 43(6) , 1123-1134.	(III)	499	2.89
225 TFA		PhSO ₂ -	H-	Intermediate 8	Intermediate 51	(III)	444	3.21
226 TFA		PhSO ₂ -	H-	Intermediate 8	3,5-difluoroaniline / Aldrich	(III)	440	3.12
227 TFA		PhSO ₂ -	H-	Intermediate 8	4-amino-2- fluorophenol / Apollo	(III)	438	2.56
228 TFA		PhSO ₂ -	H-	Intermediate 8	3,4-difluoroaniline / Aldrich	(III)	440	3.03
229 TFA		PhSO ₂ -	H-	Intermediate 8	methyl 4-amino- 2-methoxy benzoate / Avocado	(III)	492	2.89













230 HCl		PhSO ₂ -	Me-	Intermediate 16	2-(3-aminophenoxy) ethanol/ Key Organics Ltd.	(I)	478	2.22
231 HCl		PhSO ₂ -	Me-	Intermediate 16	3-aminothiophene- 2-carboxylic acid methyl ester / Avocado	(I)	482	2.79
232 HCl		PhSO ₂ -	Me-	Intermediate 16	5-amino-2- methoxyphenol / Aldrich	(I)	464	2.1
233 HCl		PhSO ₂ -	Me-	Intermediate 16	5-aminotetralin / Aldrich	(I)	472	2.64
234 HCl		PhSO ₂ -	Me-	Intermediate 16	ethyl 3- aminobenzoate / Aldrich	(I)	490	2.66
235 HCl		PhSO ₂ -	Me-	Intermediate 16	5-amino-2- methylphenol/ TCI America	(I)	448	2.31
236 HCl		PhSO ₂ -	Me-	Intermediate 16	1,1-dimethylethyl [(3-aminophenyl) methyl]carbamate / J. Med. Chem., 2003, 46 (9), 1661-1669.	(I)	547	2.64

















237 HCl		PhSO ₂ -	Me-	Intermediate 16	4-amino- <i>N</i> -methylbenzamide / Buttpark	(I)	475	2.27
238 HCl		PhSO ₂ -	Me-	Intermediate 16	3-aminobenzyl alcohol / Aldrich	(I)	448	2.13
239 TFA		PhSO ₂ -	Me-	Intermediate 16	3-aminobenzoyl methylamide / Buttpark	(III)	475	2.55
240 TFA		PhSO ₂ -	Me-	Intermediate 16	3-aminophenol / Aldrich	(III)	434	2.64
241 TFA		PhSO ₂ -	Me-	Intermediate 16	5-amino-1-ethylpyrazole / Aldrich	(III)	436	2.75
242 TFA		PhSO ₂ -	Me-	Intermediate 16	3-cyano-4-fluoroaniline hydrochloride / Combiblocks	(III)	461	3.04
243 TFA		PhSO ₂ -	Me-	Intermediate 16	1-acetyl-6-aminoindoline / Sigma	(III)	501	2.59















244 HCl		iPrSO ₂ -	H-	Intermediate 30	2,3-dihydro-1- benzofuran-4- amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(I)	412	2.36
245 HCl		iPrSO ₂ -	H-	Intermediate 30	3-amino- acetophenone/ Aldrich	(I)	412	2.33
246 HCl		iPrSO ₂ -	H-	Intermediate 30	1-methyl-1H- indazol-6-amine hydrochloride/ <i>Synth. Comm.</i> , 1996, 26(13) , 2443-2447.	(I)	424	2.22
247 HCl		iPrSO ₂ -	H-	Intermediate 30	4-fluoro-3- methoxyaniline/ Apollo-Chem	(I)	418	2.38
248 HCl		iPrSO ₂ -	H-	Intermediate 30	2,3-dihydro-1,4- benzodioxin-5- amine hydrochloride / WO 9703067 A1	(I)	428	2.30
249 HCl		iPrSO ₂ -	H-	Intermediate 30	3-chloroaniline/ Aldrich	(I)	404	2.64
250 HCl		iPrSO ₂ -	H-	Intermediate 30	3- aminobenzonitrile / Aldrich	(I)	395	2.40
251 HCl		iPrSO ₂ -	H-	Intermediate 30	3-methylaniline hydrochloride/ TCI- US	(I)	384	2.41
252 HCl		iPrSO ₂ -	H-	Intermediate 30	3-aminopyridine/ Aldrich	(I)	371	1.92












253 HCl		iPrSO ₂ -	H-	Intermediate 30	6-aminobenzo thiazole / Lancaster	(I)	427	2.20
254 HCl			H-	Intermediate 31	3-iodoaniline/ Aldrich	(I)	560	3.06
255 HCl			H-	Intermediate 31	3-amino- acetophenone / Aldrich	(I)	476	2.68
256 HCl			H-	Intermediate 31	1-methyl-1H- benzimidazol-6- amine/ <i>Heterocycles</i> , 1991, 32(5) , 1003-12.	(I)	488	2.17
257 HCl			H-	Intermediate 31	2,3-dihydro-1- benzofuran-4- amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(I)	476	2.76
258 HCl			H-	Intermediate 31	3-fluoroaniline/ Aldrich	(I)	452	2.89
259 HCl			H-	Intermediate 31	4-fluoro-3- methoxyaniline/ Apollo-Chem	(I)	482	2.78
260 HCl			H-	Intermediate 31	6-aminobenzo thiazole / Lancaster	(I)	491	2.61








261 HCl			H-	Intermediate 31	1-methyl-1H-indazol-6-amine hydrochloride/ <i>Synth. Comm.</i> , 1996, 26(13) , 2443-2447.	(I)	488	2.62
262 HCl			H-	Intermediate 31	3-aminobenzonitrile / Aldrich	(I)	459	2.86
263 HCl			H-	Intermediate 31	3-chloroaniline/ Aldrich	(I)	468	3.02
264 HCl			H-	Intermediate 31	3-methylaniline hydrochloride/ TCI- US	(I)	448	2.78
265			Me-	Intermediate 32	3-methoxyaniline/ Aldrich	(II)	478	3.04
266 TFA			Me-	Intermediate 32	2,3-dihydro-1-benzofuran-4-amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(III)	490	2.45
267 TFA			Me-	Intermediate 32	3-aminoacetophenone/ Aldrich	(III)	490	2.41
268 TFA			Me-	Intermediate 32	1-methyl-1H-indazol-6-amine hydrochloride/ <i>Synth. Comm.</i> , 1996, 26(13) ,	(III)	502	2.36









					2443-2447.			
269 TFA			Me-	Intermediate 32	2,3-dihydro-1,4-benzodioxin-5-amine hydrochloride / WO 9703067 A1	(III)	506	2.36
270 TFA			Me-	Intermediate 32	3-chloroaniline/ Aldrich	(III)	482	2.72
271 TFA			Me-	Intermediate 32	3-aminobenzonitrile / Aldrich	(III)	473	2.62
272 TFA			Me-	Intermediate 32	6-aminobenzo thiazole/ Lancaster	(III)	505	2.36
273 TFA			Me-	Intermediate 32	3-fluoroaniline/ Aldrich	(III)	466	2.6
274 HCl			H-	Intermediate 31	2-(3-aminophenoxy) ethanol hydrochloride/ <i>J. Am. Chem. Soc.</i> , 1937, 59 ; 1716	(I)	494	2.15









275 HCl			H-	Intermediate 31	3-aminophenol/ Aldrich	(I)	450	2.15
276 HCl			H-	Intermediate 31	ethyl 3-aminobenzoate/ Aldrich	(I)	506	2.51
277 HCl			H-	Intermediate 31	5-amino-2-methylphenol/ TCI America	(I)	464	2.27
278 HCl			H-	Intermediate 31	4-amino-N-methylbenzamide / Buttpark	(I)	491	2.16
279 HCl			H-	Intermediate 31	3-aminobenzyl- alcohol/ Aldrich	(I)	464	2.07
280 TFA			H-	Intermediate 31	methyl 3-aminothiophene- 2-carboxylate/ Avocado	(III)	498	2.95
281 TFA			H-	Intermediate 31	5-amino-2-methoxyphenol/ Aldrich	(III)	480	2.37
282 TFA			H-	Intermediate 31	3-amino-N-methylbenzamide / Buttpark	(III)	491	2.43









283 TFA			H-	Intermediate 31	1-amino-5,6,7,8-tetrahydro naphthalene / Aldrich	(III)	488	2.96
284 TFA			H-	Intermediate 31	4-amino-2-methoxyphenol/ WO 2003049702 A2	(III)	480	2.34
285 TFA			Me-	Intermediate 34	2,3-dihydro-1-benzofuran-4-amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(III)	474	2.53
286 TFA			Me-	Intermediate 34	3-aminoacetophenone/ Aldrich	(III)	474	2.49
287 TFA			Me-	Intermediate 34	1-methyl-1H-indazol-6-amine hydrochloride/ <i>Synth. Comm.</i> , 1996, 26(13) , 2443-2447.	(III)	486	2.41
288 TFA			Me-	Intermediate 34	2,3-dihydro-1,4-benzodioxin-5-amine hydrochloride / WO 9703067 A1	(III)	490	2.43
289 TFA			Me-	Intermediate 34	3-chloroaniline/ Aldrich	(III)	466	2.82









290 TFA			Me-	Intermediate 34	3-fluoroaniline/ Aldrich	(III)	450	2.70
291 TFA			Me-	Intermediate 34	3-aminobenzonitrile / Aldrich	(III)	457	2.71
292 TFA			Me-	Intermediate 34	3-methoxyaniline/ Aldrich	(III)	462	2.51
293 HCl		MeSO ₂ -	Me-	Intermediate 33	3-methoxyaniline/ Aldrich	(I)	386	2.20
294 HCl		MeSO ₂ -	Me-	Intermediate 33	4-fluoro-3- methoxyaniline/ Apollo-Chem	(I)	404	2.23
295 HCl		MeSO ₂ -	MeO-	Intermediate 50	3-methoxyaniline/ Aldrich	(I)	402	2.09
296 HCl		MeSO ₂ -	MeO-	Intermediate 50	4-fluoro-3- methoxyaniline/ Apollo-Chem	(I)	420	2.12
297 HCl		PhSO ₂ -	Me-	Intermediate 16	3-amino- acetophenone/ Aldrich	(I)	460	2.82

298 HCl		PhSO ₂ -	Me-	Intermediate 16	1-methyl-1H-benzimidazol-6-amine/ <i>Heterocycles</i> , 1991, 32(5) , 1003-12.	(I)	472	2.24
299 HCl		PhSO ₂ -	Me-	Intermediate 16	2,3-dihydro-1-benzofuran-4-amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(I)	460	2.90
300 HCl		PhSO ₂ -	Me-	Intermediate 16	3-fluoroaniline/ Aldrich	(I)	436	3.07
301 TFA		PhSO ₂ -	Me-	Intermediate 16	2,3-dihydro-1,4-benzodioxin-5-amine hydrochloride / WO 9703067 A1	(III)	476	2.77
302 HCl		PhSO ₂ -	Me-	Intermediate 16	6-aminobenzo thiazole / Lancaster	(I)	475	2.75
303 HCl		PhSO ₂ -	Me-	Intermediate 16	1-methyl-1H-indazol-6-amine hydrochloride / <i>Synth. Comm.</i> , 1996, 26(13) , 2443-2447.	(I)	472	2.74
304 HCl		PhSO ₂ -	Me-	Intermediate 16	3-aminobenzonitrile / Aldrich	(I)	443	3.01

305 HCl		PhSO ₂ -	Me-	Intermediate 16	3-chloroaniline/ Aldrich	(I)	452	3.21
306 HCl		PhSO ₂ -	Me-	Intermediate 16	3-methylaniline / Aldrich	(I)	432	2.93
307 HCl		MeSO ₂ -	Me-	Intermediate 33	2,3-dihydro-1- benzofuran-4- amine hydrobromide/ <i>J. Heterocyclic Chem.</i> , 1980, 17(6) , 1333-5.	(I)	398	2.38
308 TFA		MeSO ₂ -	Me-	Intermediate 33	3-aminopyridine / Aldrich	(III)	357	1.95
309 HCl		MeSO ₂ -	Me-	Intermediate 33	1-methyl-1H- indazol-6-amine hydrochloride/ <i>Synth. Comm.</i> , 1996, 26(13) , 2443-2447.	(I)	410	2.28
310 HCl		MeSO ₂ -	Me-	Intermediate 33	3-chloroaniline/ Aldrich	(I)	390	2.68
311 HCl		MeSO ₂ -	Me-	Intermediate 33	3-fluoroaniline/ Aldrich	(I)	374	2.49
312 HCl		MeSO ₂ -	Me-	Intermediate 33	3- aminobenzonitrile / Aldrich	(I)	381	2.44

313 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1-methyl-1H-benzimidazol-6-amine/ <i>Heterocycles</i> , 1991, 32(5) , 1003-12.	(I)	410	1.87
314 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	3-methylaniline/ Aldrich	(I)	370	2.40
315 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1-ethyl-1H-pyrazol-5-amine / Aldrich	(I)	374	2.22
316 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	5-(methoxy)-3-pyridinamine / <i>Australian J. Chem.</i> , 1981, 34(4) , 927-32	(I)	387	2.08
317 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	5-methyl-3-pyridinamine / Synchem	(I)	371	1.90
318 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	5-amino-2-fluorobenzonitrile / Matrix Scientific	(I)	399	2.38
329 HCl		PhSO ₂ ⁻	Me-	Intermediate 16	2-methoxybenzylamine / Aldrich	(I)	462	2.26
479 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1-methyl-1H-pyrazol-5-amine / Apollo Chem	(IV)	360	2.09

496 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1,3-benzodioxol-4-amine / <i>J. Med. Chem.</i> , 2002, 45(19) , 4128-4139	(I)	400	2.22
497 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	4-fluoro-3-methylaniline / Fluoro Chem	(I)	388	2.35
498 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	3-chloro-4-fluoroaniline / Aldrich	(IV)	408	2.59
499 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	5,6,7,8-tetrahydro-1-naphthalenamine / Fluka	(I)	410	2.57
500 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2,3-difluoroaniline / Aldrich	(IV)	392	2.52
501 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	3-chloro-2-fluoroaniline / Aldrich	(I)	408	2.67
502 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	3,5-difluoroaniline / Aldrich	(I)	392	2.60
503 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1,3-benzothiazol-6-amine / Maybridge	(I)	413	2.13

504 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	3,4-difluoroaniline / Aldrich	(I)	392	2.43
505 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2,4-difluoroaniline / Aldrich	(I)	392	2.33
508 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1-methyl-2,3-dihydro-1H-indol-4-amine / Intermediate 71	(IV)	411	2.2
509 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	1-methyl-1H-indazol-4-amine / <i>J. Med. Chem.</i> , 2002, 45 (3), 740-743	(I)	410	2.21
510 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2,3-dihydro-1-benzofuran-7-amine / WO9517401 A1	(I)	398	2.15
511 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2,3-dihydro-1H-inden-4-amine / Aldrich	(I)	396	2.46
512 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	4-amino-2,3-dihydro-1H-inden-1-one / Davos	(IV)	410	2.13
520 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2-methyl-4-pyridinamine/ Asym Chem	(I)	371	1.7

542 HCl		MeSO ₂ -	Me-	Intermediate 33	1,3-dihydro-2-benzofuran-4-amine / US 4521241	(I)	398	2.16
587 HCl		MeSO ₂ -	Me-	Intermediate 33	3,4-dihydro-2H-chromen-5-ylamine / J. Heterocyclic Chem., 1973, 10(4), 623-9	(I)	412	2.26


(a) Salt forms: HCl = hydrochloride
TFA = trifluoroacetate

(b) Isolation Method:

- 5 (I) Filtered off directly from the reaction mixture.
(II) Mass Directed preparative HPLC Method A.
(III) Mass Directed preparative HPLC Method B.
(IV) Mass Directed preparative HPLC Method C.

10


Example 14. 4-[(3-Chlorophenyl)(methyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide



- 15 Intermediate 9 (0.023g) was dissolved in 1-methyl-2-pyrrolidinone (1ml), and 3-chloro-N-methylaniline (available from Avocado) (0.012ml) was added. The mixture was stirred under microwave irradiation (power 150W) for 10min at 180°C and for a further 10min (power 150W) at 150°C. Purification by mass directed HPLC (Method A) gave the title compound (0.015g).

LC/MS R_t 2.58min m/z 390 $[MH^+]$


Example 112. 4-({[2-(Methyloxy)phenyl]methyl}amino)-6-(phenylsulfonyl)-3-quinolinecarboxamide











Intermediate 8 (0.017g) was taken up in acetonitrile (1.5ml) to give a slurry. 2-Methoxy benzylamine (available from Aldrich) (0.021g) and *N,N*-diisopropylethylamine (0.050ml) were added and the resultant mixture was heated under reflux for 16h. The mixture was cooled, the solvent evaporated *in vacuo* and the residue purified by mass directed HPLC (Method A) to give the title compound (0.014g).











LC/MS R_t 2.73min m/z 448 $[MH^+]$











Similarly prepared were the following:



Ex. No (a)	R^1NH	R^3SO_2	Starting Material	Amine reagent/ Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
---------------	---------	-----------	-------------------	--------------------------	-------------------------	-------------	---------------------

101 (c) HCl		MeSO ₂ -	Intermediate 9	1-phenylmethanamine/ Aldrich	(I)	356	2.07
102		MeSO ₂ -	Intermediate 9	tetrahydro-2H-pyran-3-amine hydrochloride/ <i>Anales de Quimica, Serie C: Quimica Organica y Bioquimica</i> , 1988, 84(2) , 148-55.	(II)	350	1.78
103		PhSO ₂ -	Intermediate 8	2-(aminomethyl)phenol/ Buttpark	(I)	434	2.56
104		PhSO ₂ -	Intermediate 8	2,3-dihydro-1H-inden-2-amine hydrochloride / Aldrich	(I)	444	2.85
105		PhSO ₂ -	Intermediate 8	cyclopropylamine/ Aldrich	(I)	368	2.37
106		PhSO ₂ -	Intermediate 8	[4-(aminomethyl)phenyl] dimethylamine hydrochloride/ Aldrich	(I)	461	2.69
107		PhSO ₂ -	Intermediate 8	N-[3-(aminomethyl)phenyl] methanesulfonamide trifluoroacetate/ <i>J. Med. Chem.</i> , 1999, 42(14) , 2504-2526.	(I)	511	2.52
108		PhSO ₂ -	Intermediate 8	2-aminocyclohexanol/ TCI-US	(I)	426	2.49

109		PhSO ₂ ⁻	Intermediate 8	[(1-methyl-1H-pyrazol-4-yl)methyl]amine/ Zelinsky-BB; CA 400877-05-6	(I)	422	2.28
110		PhSO ₂ ⁻	Intermediate 8	(2-pyridinyl)methyl amine hydrochloride/ Aldrich	(I)	419	2.35
111		PhSO ₂ ⁻	Intermediate 8	1,2,3,4-tetrahydro-1-naphthalenamine hydrochloride/ Aldrich	(I)	458	2.92
113		PhSO ₂ ⁻	Intermediate 8	(cyclohexylmethyl) amine / Aldrich	(I)	424	2.87
114		PhSO ₂ ⁻	Intermediate 8	{[4-(methoxy)phenyl] methyl}amine hydrochloride/ Aldrich	(I)	448	2.69
115		PhSO ₂ ⁻	Intermediate 8	(phenylmethyl)amine hydrochloride/ Aldrich	(I)	418	2.67
116		PhSO ₂ ⁻	Intermediate 8	cyclohexylamine hydrochloride/ Acros	(II)	410	2.59
117		PhSO ₂ ⁻	Intermediate 8	2-methyl-1-propanamine trifluoroacetate/ Aldrich	(II)	384	2.43
118		PhSO ₂ ⁻	Intermediate 8	[2-(3-pyridinyl)ethyl] amine / Lancaster	(II)	433	2.08
119		PhSO ₂ ⁻	Intermediate 8	[2-(4-pyridinyl)ethyl] amine / Maybridge	(II)	433	2.01

120		PhSO ₂ ⁻	Intermediate 8	2-phenylethanamine/ Aldrich	(II)	432	2.68
121		PhSO ₂ ⁻	Intermediate 8	(3-pyridinylmethyl) amine / Aldrich	(II)	419	2.12
122		PhSO ₂ ⁻	Intermediate 8	{[3,5- bis(methoxy)phenyl] methyl}amine hydrochloride/ Aldrich	(II)	478	2.68
123		PhSO ₂ ⁻	Intermediate 8	tetrahydro-2H-pyran- 3-amine hydrochloride/ <i>Anales de Quimica, Serie C: Quimica Organica y Bioquimica</i> , 1988, 84(2) , 148-55.	(II)	412	2.20
124		PhSO ₂ ⁻	Intermediate 8	4-amino cyclohexanone/ <i>Nouveau Journal de Chimie</i> , 1984, 8(7) , 459-67.	(II)	424	2.15
125 (c) HCl		PhSO ₂ ⁻	Intermediate 8	{[3-chloro-4- (methoxy)phenyl] methyl}amine/ Apin Chemicals	(I)	482	2.56
126			Intermediate 7	cyclohexylamine/ Aldrich	(II)	402	2.49
127			Intermediate 7	2-methyl-1- propanamine/ Aldrich	(II)	376	2.34

(a) Salt forms: HCl = hydrochloride


(b) Isolation Method:

(I) Filtered off directly from the reaction mixture; it is thought that compounds isolated by this method are free bases, apart from Examples 101 and 125 which are thought to be hydrochloride salts.

(II) Mass Directed preparative HPLC Method A.

5 (c) No *N,N*-diisopropylethylamine was used in the preparation of Examples 101 and 125.

The following were made in a similar manner to Example 112:



10

Ex. No. (a)	R^1R^2N-	R^3SO_2-	$R^{20}-$	Starting Material	Amine Reagent/ Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
319		PhSO ₂ -	H-	Intermediate 8	<i>N</i> ,3-dimethylaniline / Acros	(II)	432	2.97
320		PhSO ₂ -	H-	Intermediate 8	3-chloro- <i>N</i> -methylaniline / Maybridge	(II)	452	3.00
321 TFA		PhSO ₂ -	H-	Intermediate 8	3-aminoquinuclidine dihydrochloride / Aldrich	(III)	437	1.98
330 TFA		PhSO ₂ -	Me-	Intermediate 16	(3-pyridinylmethyl) amine / Aldrich	(III)	433	2.08

331 HCl			H-	Intermediate 31	2-(aminomethyl) phenol/ Buttpark	(I*)	464	2.14
332 TFA			H-	Intermediate 31	(3-pyridinylmethyl) amine/ Aldrich	(III)	449	2.06
333 TFA			H-	Intermediate 31	[2-(4-pyridinyl)ethyl] amine/ Maybridge	(III)	463	1.93
334 HCl		MeSO ₂ ⁻	Me-	Intermediate 33	2-methyl-2- propanamine / Aldrich	(IV)	336	2.04

(a) Salt forms: HCl = hydrochloride

TFA = trifluoroacetate

(b) Isolation Method:

- 5 (I) Filtered off directly from the reaction mixture; it is thought that compounds isolated by this method are free bases.


(I*) No base is used in the reaction procedure . Filtered off directly from the reaction mixture; it is thought that compounds isolated by this method are hydrochloride salts.

- 10 (II) Mass Directed HPLC Method A; it is thought that compounds isolated by this method are free bases unless the R¹ or R³ groups contain basic moieties, in which case formate salts may be formed.

(III) Mass Directed HPLC Method B; it is thought that compounds isolated by this method are trifluoroacetate salts.

- 15 (IV) Mass Directed HPLC Method C; it is thought that compounds isolated by this method are hydrochloride salts.

Example 133. 6-[(1,1-Dimethylethyl)thio]-4-[[3-(methoxy)phenyl]amino]-3-quinolinecarboxamide



5


Intermediate 14 (0.050g), potassium *tert*-butoxide (0.015g) and *tert*-butylmercaptan (0.0135ml) were added to a stirred solution of tris(dibenzylideneacetone)dipalladium (0) (0.007g) and (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.005g) in toluene (2ml), and the mixture was heated at 100°C for 3.5h and left to cool. The solvent was evaporated *in vacuo* to leave a brown solid (0.072g), which was purified by Mass Directed Preparative HPLC (Method A) to give the title compound (0.008g).

10











LC/MS R_t 2.83min m/z 382 $[MH^+]$.

Similarly prepared from Intermediate 14 were the following:

15



Ex. No.	R ³ S-	Thiol reagent/ Source	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
132		cyclohexanethiol / Aldrich	(II)	408	2.99

135		N-(2-mercaptoethyl) acetamide / Aldrich	(II)	411	2.11
136		2,6-dichlorobenzene thiol / Aldrich	(II)	471	2.77
137		2-methyl-1- propanethiol / Aldrich	(II)	382	2.66
138		1,3-oxazole-2(3H)- thione/ <i>Can. J. Chem.</i> , 1972, 50(18), 3082-3.	(II)	393	2.29
139		5-methyl-1,3,4- oxadiazole-2(3H)- thione/ US 5670526 A	(II)	408	2.22
140		phenylmethane thiol/ Aldrich	(II)	416	2.66
141		4-fluorobenzene thiol/ Aldrich	(IV)	420	2.87
142		4-(methoxy) benzenethiol/ Aldrich	(II)	432	2.90
143		cyclopentanethiol/ Aldrich	(I)	394	2.98
144		benzenethiol/ Aldrich	(I)	402	2.96

(b) Isolation Method:


(I) Filtered off directly from the reaction mixture; it is thought that compounds isolated by this method are free bases.

(II) Mass Directed preparative HPLC Method A.

(IV) Purified by chromatography on silica gel, eluting with dichloromethane followed by ethyl acetate. It is thought that compounds isolated by this method are free bases.



















5

The following were prepared in a similar manner to Example 133, using *N,N*-dimethylformamide as the reaction solvent:











































10



















Ex. No.	R ¹ NH-	R ³ S-	R ²⁰ -	Starting Material	Thiol Reagent / Source	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
337			Me-	Intermediate 35	2-mercaptoethanol/ Sigma	(I)	402	2.19
338			Me-	Intermediate 35	1,2,4-triazole-3-thiol/ Aldrich	(I)	425	2.12
339			Me-	Intermediate 35	1-methyl-2-mercaptoimidazole / Aldrich	(I)	438	2.15
340			Me-	Intermediate 35	2-mercaptoimidazole / Aldrich	(I)	424	1.96
341			Me-	Intermediate 35	2-benzimidazolethiol/ Aldrich	(I)	474	2.54

342			Me-	Intermediate 35	4-methyl-1,3-oxazole-2(3H)-thione <i>J. Org. Chem.</i> , 1967, 32(7) , 2079-81.	(I)	439	2.67
343			Me-	Intermediate 35	2-furanyl methanethiol / Aldrich	(I)	438	2.79
344			Me-	Intermediate 36	1,1-dimethylethyl (2-mercaptoethyl) carbamate/ Aldrich	(V)	495	2.73
345			Me-	Intermediate 36	1,1-dimethylethyl 4-mercapto-1- piperidinecarboxyla te/ US 5317025 A	(V)	535	3.22
563			Me-	Intermediate 62	tetrahydro-3- furanthiol/ <i>Advan.</i> <i>Carbohydrate</i> <i>Chem.</i> (1963), 18 123-99	(I)	381	2.17
564			Me-	Intermediate 61	tetrahydro-3- furanthiol / <i>Advan.</i> <i>Carbohydrate</i> <i>Chem.</i> (1963), 18 123-99	(I)	398	2.65
565			Me-	Intermediate 61	tetrahydro-2H- pyran-4-thiol / WO98/05635	(I)	412	2.70
566			Me-	Intermediate 62	tetrahydro-2H- pyran-4-thiol/ WO98/05635	(I)	395	2.20
569			Me-	Intermediate 62	1,1-dimethylethyl 4-mercapto-1- piperidine carboxylate /US 5317025 A	(I)	494	2.80

570			Me-	Intermediate 61	N-(2-mercaptoethyl) acetamide / Aldrich	(I)	413	2.20
572			Me-	Intermediate 62	N-(2-mercaptoethyl)acet amide / Aldrich	(V)	396	1.90
516			Me-	Intermediate 35	1,1-dimethylethyl 4-mercapto-1- piperidine carboxylate/ US5317025A	(V)	541	3.10
517			Me-	Intermediate 35	1,1-dimethylethyl (2-mercaptoethyl) carbamate/ Aldrich	(V)	501	3.68
528			Me-	Intermediate 36	2-mercaptoethanol / Aldrich	(I)	396	2.20
603			Me-	Intermediate 61	1,1-dimethylethyl (2-mercaptoethyl) carbamate	(I)	471	2.80
634			Me-	Intermediate 62	ethanethiol / Aldrich	(I)	339	2.32
635			Me-	Intermediate 62	1-propanethiol / Aldrich	(I)	353	2.71
636			Me-	Intermediate 62	2-propanethiol / Aldrich	(I)	353	2.47
637			Me-	Intermediate 62	2-methyl-2- propanethiol / Aldrich	(I)	367	2.58

638			Cl-	Intermediate 96	ethanethiol / Aldrich	(I)	359	2.59
639			Cl-	Intermediate 96	1-propanethiol / Aldrich	(I)	373	2.76
640			Cl-	Intermediate 96	2-propanethiol / Aldrich	(I)	373	2.71
641			Cl-	Intermediate 96	2-methyl-2- propanethiol / Aldrich	(I)	387	2.93
642			Me-	Intermediate 62	ethanethiol / Aldrich	(I)	373	2.81
643			Me-	Intermediate 97	1-propanethiol / Aldrich	(I)	387	3.06
644			Me-	Intermediate 97	2-propanethiol / Aldrich	(I)	387	2.95
645			Me-	Intermediate 97	2-methyl-2- propanethiol / Aldrich	(I)	401	3.16
646			Cl-	Intermediate 100	ethanethiol / Aldrich	(I)	393	3.1
647			Cl-	Intermediate 100	1-propanethiol / Aldrich	(I)	407	3.3

648			Cl-	Intermediate 100	2-propanethiol / Aldrich	(I)	407	3.25
649			Cl-	Intermediate 100	2-methyl-2- propanethiol / Aldrich	(I)	421	3.33
650			Me-	Intermediate 98	ethanethiol / Aldrich	(I)	357	2.59
651			Me-	Intermediate 98	1-propanethiol / Aldrich	(I)	371	2.84
652			Me-	Intermediate 98	2-propanethiol / Aldrich	(I)	371	2.84
653			Me-	Intermediate 98	2-methyl-2- propanethiol / Aldrich	(I)	385	2.94
654			Cl-	Intermediate 99	ethanethiol / Aldrich	(I)	377	2.88
655			Cl-	Intermediate 99	1-propanethiol / Aldrich	(I)	391	3.07
656			Cl-	Intermediate 99	2-propanethiol / Aldrich	(I)	391	3.03
657			Cl-	Intermediate 99	2-methyl-2- propanethiol / Aldrich	(I)	405	3.14


658			Cl-	Intermediate 102	ethanethiol / Aldrich	(I)	376	2.94
659			Cl-	Intermediate 102	1-propanethiol / Aldrich	(I)	390	3.08
660			Cl-	Intermediate 102	2-propanethiol / Aldrich	(I)	390	3.02
661			Cl-	Intermediate 102	2-methyl-2-propanethiol / Aldrich	(I)	404	3.21
662			Me-	Intermediate 101	ethanethiol / Aldrich	(I)	356	2.5
663			Me-	Intermediate 101	1-propanethiol / Aldrich	(I)	370	2.65
664			Me-	Intermediate 101	2-propanethiol / Aldrich	(I)	370	2.61
665			Me-	Intermediate 101	2-methyl-2-propanethiol / Aldrich	(I)	384	2.89
681			Me-	Intermediate 61	N-(2-mercaptoethyl)-N-methylacetamide / Tetrahedron 1986, 42 (5), 1449	(I)	427	2.30

(b) Isolation Method:

(I) These were purified by SCX column, eluting with ammonia/ methanol.


(V) These were purified by chromatography on silica gel (eluting with ethyl acetate/cyclohexane) followed by trituration with cyclohexane to give the pure product; it is thought that compounds isolated by this method are free bases.












5 **Example 577:** 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylthio)-3-quinolinecarboxamide.



















A stirred mixture of Intermediate 36 (0.2g), sodium thiomethoxide (0.058g),
 tris(dibenzylideneacetone)dipalladium(0) (0.076g), (oxidi-2,1-phenylene)-
 10 bis(diphenylphosphine) (0.045g), and potassium *tert*-butoxide (0.047g) in *N,N*-
 dimethylformamide (10ml) was heated at 100°C under nitrogen for 18h. The cooled
 reaction mixture was applied directly to an SCX cartridge (10g) and eluted with methanol
 (150ml) followed by 2M ammonia in methanol (100ml). Evaporation of the methanol/
 ammonia fraction gave the title compound as a yellow solid (0.13g).
 15 LC/MS R_t 2.48min, m/z 366 $[MH^+]$

The following were prepared in a similar manner to Example 577, but without adding
 potassium *tert*-butoxide to the reaction mixture:




Ex. No.	R ¹ NH-	R ²⁰ -	Starting Material	LCMS MH ⁺	LCMS R _t (min)
577		Me-	Intermediate 36	382	2.00
604		Et-	Intermediate 74	386	2.57
605		Et-	Intermediate 75	356	2.76
606		Et-	Intermediate 76	372	2.95
607		Et-	Intermediate 77	363	2.71
608		Et-	Intermediate 78	352	2.65
609		Et-	Intermediate 79	392	2.44
610		Et-	Intermediate 80	380	2.62
611		Et-	Intermediate 81	339	2.2
612		Et-	Intermediate 82	357	2.55
613		Et-	Intermediate 83	373	2.73

614		F-	Intermediate 85	363	2.69
615		F-	Intermediate 86	329	2.16
616		F-	Intermediate 84	347	2.53
617		F-	Intermediate 89	346	2.87
618		F-	Intermediate 90	363	3.06
619		F-	Intermediate 91	342	2.91
620		F-	Intermediate 92	353	2.77
621		F-	Intermediate 93	382	2.56
622		F-	Intermediate 94	376	2.88
623		F-	Intermediate 88	370	2.81
682		Cl-	Intermediate 38	358	3.27
683		Cl-	Intermediate 39	392	3.06

684		Cl-	Intermediate 40	386	3.10
685		Cl-	Intermediate 41	369	3.01
686		Cl-	Intermediate 42	362	3.20
687		Cl-	Intermediate 43	398	2.82

Example 134. 4-([3-(Methyloxy)phenyl]amino)-6-([4-(methyloxy)phenyl]methyl)thio-3-quinolinecarboxamide




5

Intermediate 14 (0.020g), potassium *tert*-butoxide (0.0061g) and [4-(methyloxy)phenyl]methanethiol (available from Aldrich) (0.007ml) were added to a stirred solution of tris(dibenzylideneacetone)dipalladium(0) (0.002g) and (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.002g) in *N,N*-dimethylformamide (1.5ml), and the mixture was heated under microwave irradiation at 60°C for 8min. The solvent was evaporated *in vacuo*, and the residue purified by Mass Directed Preparative HPLC (Method A) to give the title compound (0.0048g).

LC/MS R_t 2.93min m/z 446 [MH^+]

15

Example 129. 6-[(1,1-Dimethylethyl)sulfonyl]-4-[[3-(methoxy)phenyl]amino]-3-quinolinecarboxamide




5

Example 133 (0.010g) was dissolved in *N,N*-dimethylformamide (2ml) and anisole (0.013ml) was added. Oxone (0.075g) was added and the mixture stirred for 16h at room temperature. After quenching with 1M aqueous sodium sulphite, the mixture was extracted with dichloromethane; the organic layer was dried over magnesium sulphate and evaporated *in vacuo* to give a yellow solid. Purification by mass directed HPLC (Method A) gave the title compound (0.005g).

LC/MS R_t 2.48min m/z 414 $[MH^+]$.

15 Similarly prepared were the following:




Ex. No.	R^3SO_2-	Starting Material	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
---------	------------	-------------------	----------------------	-------------	------------------

128		Example 142	(II)	464.09	2.83
130 (a)		Example 135	(II)	443.1	2.11















(a) No anisole was used in the reaction mixture in this Example.

(b) Isolation Method: (II) Mass Directed preparative HPLC (Method A).

- 5 The following were prepared in a similar manner to Example 129, but without the addition of anisole to the reaction mixture:



Ex. No. (a)	R ¹ NH-	R ³ SO ₂ -	R ²⁰ -	Starting Material	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
346 HCl			Me-	Example 337	(IV)	434	2.10
347 HCl			Me-	Example 338	(IV)	457	2.30
348 HCl			Me-	Example 339	(IV)	470	2.46

349 HCl			Me-	Example 340	(IV)	456	2.27
350 HCl			Me-	Example 341	(IV)	506	2.79
351 HCl			Me-	Example 343	(IV)	470	2.66
352			Me-	Example 345	(V)	567	3.08
353			Me-	Example 344	(VI)	527.5	2.76
369 HCl		MeSO ₂ ⁻	Cl-	Example 682	(IV)	390	2.77
370 HCl		MeSO ₂ ⁻	Cl-	Example 683	(IV)	424	2.63
371 HCl		MeSO ₂ ⁻	Cl-	Example 684	(IV)	418	2.77
372 HCl		MeSO ₂ ⁻	Cl-	Example 685	(IV)	401	2.57

373 HCl		MeSO ₂ ⁻	Cl ⁻	Example 686	(IV)	394	2.70
374 HCl		MeSO ₂ ⁻	Cl ⁻	Example 687	(IV)	430	2.51
478			Me ⁻	Intermediate 29	(VI)	519	2.55
602			Me ⁻	Example 603	(VII)	503	2.90
515			Me ⁻	Example 562	(VI)	449	2.60

(a) Salt forms: HCl = hydrochloride.

(b) Isolation Method:


(IV) Mass Directed preparative HPLC (Method C).

5 (V) Column chromatography on silica gel.










(VI) Aqueous work-up.







(VII) Trituration from acetonitrile.

The following were prepared in a similar manner to Example 129 without the addition of
 10 anisole to the reaction mixture:



Ex. No. (a)	R ¹ NH-	R ²⁰ -	R ³ SO ₂ -	Starting Material	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
522		Me-		Example 521	(II)	448	2.33
591 HCl		Et-	MeSO ₂ -	Example 604	(IV)	418	2.39
592 HCl		Et-	MeSO ₂ -	Example 605	(IV)	388	2.51
593 HCl		Et-	MeSO ₂ -	Example 607	(IV)	395	2.46
594 HCl		Et-	MeSO ₂ -	Example 612	(IV)	389	2.32
595 HCl		Et-	MeSO ₂ -	Example 613	(IV)	405	2.47

596 HCl		Et-	MeSO ₂ -	Example 608	(IV)	384	2.41
597 HCl		Et-	MeSO ₂ -	Example 606	(IV)	404	2.70
598 HCl		Et-	MeSO ₂ -	Example 609	(IV)	424	2.34
599 HCl		Et-	MeSO ₂ -	Example 610	(IV)	412	2.44
600 HCl		Et-	MeSO ₂ -	Example 611	(IV)	370	1.99
624		F-	MeSO ₂ -	Example 623	(II)	402	2.52
625		F-	MeSO ₂ -	Example 617	(II)	378	2.54
626		F-	MeSO ₂ -	Example 618	(II)	395	2.70
627		F-	MeSO ₂ -	Example 619	(II)	374	2.61

628		F-	MeSO ₂ -	Example 620	(II)	385	2.42
629		F-	MeSO ₂ -	Example 621	(II)	414	2.27
630		F-	MeSO ₂ -	Example 622	(II)	408	2.45
631		F-	MeSO ₂ -	Example 615	(II)	361	1.91
632		F-	MeSO ₂ -	Example 614	(II)	395	2.32
633		F-	MeSO ₂ -	Example 616	(II)	379	2.09


(a) Salt forms: HCl = hydrochloride.

(b) Isolation method:

(II) Mass Directed preparative HPLC (Method A).

5 (IV) Mass Directed preparative HPLC (Method C).

Example 184. 8-Methyl-4-[[3-(methoxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide hydrochloride




Intermediate 16 (0.036g) was suspended in acetonitrile (2ml), 3-methoxyaniline (available from Aldrich) (0.012g) was added, and the mixture was heated under reflux for 16h. After cooling to room temperature the mixture was filtered and the residue dried to give the

5 title compound as a beige solid (0.020g).

LC/MS R_t 2.86min m/z 448 $[MH^+]$

Similarly prepared was:




10

Ex. No.	R^1NH-	R^3SO_2-	Starting Material	Amine Reagent/ Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
(a)							
185 HCl		PhSO ₂ -	Intermediate 16	4-fluoro-3-methoxyaniline / Apollo-Chem	(I)	466	2.89

(b) Isolation method: (I) Filtered off from the reaction mixture.

Example 186. 7-Methyl-4-([3-(methoxy)phenyl]amino)-6-(methylsulfonyl)-3-quinolinecarboxamide hydrochloride


15



Intermediate 17 (0.058g) was suspended in acetonitrile (2ml), 3-methoxyaniline (0.024g) (available from Aldrich) was added, and the mixture was heated under reflux for 4h. After cooling to room temperature the mixture was filtered and the residue dried to give the title compound as a beige solid (0.042g).

LC/MS R_t 2.21min m/z 386 $[MH^+]$

10 **Example 335.** 4-[(3-Aminophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide trifluoroacetate




15 To a stirred mixture of Example 187 (0.130g) in dichloromethane (5ml) was added trifluoroacetic acid (1ml). The mixture was stirred at 20°C for 1h and then the solvent was removed *in vacuo* to give the title compound as a yellow gum (0.100g).

LC/MS R_t 1.87min m/z 357 $[MH^+]$

20 Similarly prepared from example 188 was:


Example 336. 4-([3-(Aminomethyl)phenyl]amino)-6-(methylsulfonyl)-3-quinolinecarboxamide trifluoroacetate



5

LC/MS R_t 1.65min m/z 371 [MH^+]

Example 376: 1,1-Dimethylethyl 4-({3-(aminocarbonyl)-8-methyl-4-[(1-methyl-1*H*-benzimidazol-6-yl)amino]-6-quinoliny]sulfonyl)-1-piperidinecarboxylate



- 15 A mixture containing Intermediate 44 (0.500g), 1,1-dimethylethyl 4-mercapto-1-piperidinecarboxylate (0.442g, synthesised according to US5317025A), potassium *tert*-butoxide (0.248g), tris(dibenzylideneacetone) dipalladium (0.093g) and (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.091g) was dissolved in *N,N*-dimethylformamide (20ml) and stirred under an atmosphere of nitrogen at 100°C for 2h. The solvents were
- 20 concentrated *in vacuo* and the residue partitioned between ethyl acetate (100ml) and


water (100ml). The organic extract was washed with sodium bicarbonate solution followed by brine, dried over magnesium sulfate and concentrated *in vacuo* to an orange solid. This was purified by flash chromatography on silica gel eluting with a gradient of ethanol (0% to 10%) in ethyl acetate, to give the intermediate sulphide 1,1-dimethylethyl 4-({3-(aminocarbonyl)-8-methyl-4-[(1-methyl-1*H*-benzimidazol-6-yl)amino]-6-quinoliny]thio)-1-piperidinecarboxylate as a yellow solid (0.375g).

LC/MS R_t 2.63 min, m/z 547 [MH^+]

Oxone (1.6g) was added to a solution of the sulphide (0.370g) in *N,N*-dimethylformamide (10ml). The mixture was stirred at room temperature for 1h and was quenched with a solution of sodium sulphite (4g) in water (150ml). The mixture was extracted with ethyl acetate (2 x 100ml) and the combined organic suspension washed with water (2 x 100ml) and extracted *in vacuo* to a pale yellow solid. This was purified by recrystallisation from boiling methanol to give the title compound as a pale yellow powder (0.265g).

LC/MS R_t 2.47 min, m/z 579 [MH^+]

Similarly prepared were the following:



20

Ex. No.	R¹NH-	R³SO₂-	R²⁰-	Starting Material	Thiol Reagent/ Source	Isolation Method (b)	Solvent	LCMS MH⁺	LCMS R _t (min)
354			H-	Intermediate 37	Methyl 4-mercaptobenzoate/ Toronto Research Chemicals	(II)	Toluene	492	2.76

355			H-	Intermediate 37	Methyl 3-mercaptobenzoate/ Toronto Research Chemicals	(II)	Toluene	492	2.76
356			H-	Intermediate 37	3-methylbenzene thiol / Aldrich	(VII)	Toluene	448	2.90
357 HCl			H-	Intermediate 14	3,4-dimethylthio phenol/ Aldrich	(IV)	Toluene	462	3.02
358			H-	Intermediate 14	3-fluorobenzene thiol/ Avocado	(V)	Toluene	452	2.83
359 HCl			H-	Intermediate 14	4-(trifluoromethyl) thiophenol/ Fluorochem	(IV)	Toluene	502	3.16
361			H-	Intermediate 37	3-chlorothiophenol/ Aldrich	(II)	Toluene	468	2.96
362			H-	Intermediate 37	4-tert-butylthio phenol / Lancaster	(II)	Toluene	490	3.26
363			H-	Intermediate 37	3,5-dimethyl benzenethiol/ Aldrich	(II)	Toluene	462	3.05
364			H-	Intermediate 37	1,1-dimethylethyl (2-mercaptoethyl) carbamate/ Aldrich	(V)	Toluene	469	2.59

365 HCl			H-	Intermediate 37	[4-(methoxyphenyl) methanethiol / Aldrich	(IV)	Toluene	478	2.63
366			H-	Intermediate 37	4-bromothiophenol/ Aldrich	(VI)	Toluene	513	3.02
367 HCl			H-	Intermediate 37	2-mercaptoanisole/ Lancaster	(IV)	Toluene	464	2.52
368 HCl			H-	Intermediate 37	(4-chlorophenyl) methanethiol/ Aldrich	(IV)	Toluene	482	2.91
375 HCl			H-	Intermediate 37	3-methoxybenzene thiol / Aldrich	(VIII)	N,N- dimethyl- formamide	464	2.76
377			Me-	Intermediate 45	1,1-dimethylethyl 4-mercapto-1- piperidine carboxylate / US 5317025 A	(V)	N,N- dimethyl- formamide	555	2.97
378			Me-	Intermediate 46	1,1-dimethylethyl 4-mercapto-1- piperidine carboxylate / US 5317025 A	(V)	N,N- dimethyl- formamide	550	3.01
472			H-	Intermediate 14	1,1-dimethylethyl 4-mercapto-1- piperidine carboxylate / US 5317025 A	(V)	N,N- dimethyl- formamide	541	2.89

(a) Salt forms: HCl = hydrochloride

(b) Isolation Method:

(II) Mass Directed preparative HPLC (Method A).

(IV) Mass Directed preparative HPLC (Method C).


(V) Column chromatography on silica gel. Compounds isolated by this method are free bases.

(VI) Aqueous workup only with no further chromatography. Compounds isolated by this method are free bases.

(VII) Recrystallised from methanol

(VIII) Mass Directed preparative HPLC (Method A), followed by treatment with 2M HCl in ethanol.


Example 360. 6-[(4-Hydroxyphenyl)sulfonyl]-4-[(3-methoxyphenyl)amino]quinoline-3-carboxamide



Oxone (3.9g) was added to a stirred solution of Intermediate 47 (1.1g) in dry *N,N*-dimethylformamide (30ml) at room temperature for 18h. The mixture was poured into aqueous sodium sulphite solution (200ml) and extracted with ethyl acetate (3x100ml). The organic extracts were washed with water (2x100ml), dried (Na_2SO_4) and concentrated. A solution of the residual oil in tetrahydrofuran (20ml) was stirred with a 1M solution of tetrabutylammonium fluoride in tetrahydrofuran (4ml) for 1h. The solvent was removed *in vacuo* and the residue was partitioned between ethyl acetate (2x25ml) and water (2x50ml). The organic extracts were dried (Na_2SO_4) and concentrated to give the title compound as a yellow solid (0.67g).

LC/MS R_t 2.58 min, m/z 450 [MH^+]

Example 379. Methyl 3-[(3-(aminocarbonyl)-8-methyl-4-{[3-(methyloxy)phenyl]amino}-6-quinolinyl)sulfonyl]benzoate




To Intermediate 45 (0.47g) in dimethoxyethane (10ml) was added methyl 3-mercaptobenzoate (0.34ml), potassium phosphate (0.42g), copper (I) iodide (0.028g) and *N,N*-diethylsalicylamide (0.39g). The mixture was heated at 85°C for 4h before adding further methyl 3-mercaptobenzoate (0.34ml) and copper (I) iodide (0.028g). After a further 16h the reaction mixture was concentrated *in vacuo* and partitioned between ethyl acetate (150ml) and water (150ml). The organic layers were washed with brine (100ml), dried over sodium sulfate and concentrated *in vacuo* to yield a crude product which was triturated with diethyl ether (20ml). The solid obtained was collected by filtration, washed with diethyl ether (2 X 10ml) to give methyl 3-[(3-(aminocarbonyl)-8-methyl-4-[[3-(methoxy)phenyl]amino]-6-quinolinyl)thio]benzoate as a beige solid (0.37g).

LC/MS R_t 3.09min m/z 474 $[MH^+]$

To a solution of the methyl 3-[(3-(aminocarbonyl)-8-methyl-4-[[3-(methoxy)phenyl]amino]-6-quinolinyl)thio]benzoate (0.367g) in *N,N*-dimethylformamide (10ml) was added oxone (1.91g). The mixture was stirred at room temperature for 18h before quenching with aqueous sodium sulphite solution and extracting with chloroform (3 x 200ml). The organic layers were combined, washed with brine, dried over magnesium sulfate and concentrated *in vacuo* and purified by chromatography on silica gel, eluting with 2:1 ethyl acetate: cyclohexane, to give the title compound as a yellow solid (0.100g).

LC/MS R_t 3.03min m/z 506 $[MH^+]$

The following were synthesised in the same manner as Example 379, however potassium carbonate was used as a base instead of potassium phosphate and no *N,N*-diethylsalicylamide was added.



5

Ex. No. (a)	R ³ SO ₂ -	R ²⁰ -	Starting Material	Thiol Reagent/ Source	Purification Method (b)	LCMS MH ⁺	LCMS R _t (min)
380		Me-	Intermediate 45	3,4-dimethoxythiophenol Aldrich	(V)	508	2.81
381 HCl		Me-	Intermediate 45	3,4,5-tris(methoxy) benzenethiol <i>J. Am. Chem. Soc.</i> , 2002, 124 (17), 4642-4646	(IV)	538	2.92
382 HCl		H-	Intermediate 14	3,4-dimethoxythiophenol/ Avocado	(IV)	494	2.59
383 HCl		Me-	Intermediate 45	3-ethoxythiophenol/ Aldrich	(IV)	492	3.08


(a) Salt forms: HCl = hydrochloride

(b) Isolation Method:

(IV) Mass Directed preparative HPLC (Method C).

(V) Column chromatography on silica gel; it is thought that compounds isolated by this method are free bases.

5 **Example 386.** 6-(Ethylsulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide hydrochloride




Intermediate 37 (0.100g) was combined with (oxydi-2,1-phenylene)bis(diphenylphosphine) (0.011g), potassium *tert*-butoxide (0.025g) and
 10 tris(dibenzylideneacetone) dipalladium(o) (0.008g) in 1,4-dioxane (1ml). Ethanethiol (available from Aldrich, 0.023ml) was added and the mixture was stirred under microwave irradiation (power 40W) for 8 min at 90°C. The reaction was quenched by addition of 4M HCl in dioxane, then partitioned between ethyl acetate and sodium bicarbonate solution. The organic layer was concentrated *in vacuo* to give 0.090g of crude 6-(ethylthio)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide.
 15


The crude sulphide was dissolved in *N,N*-dimethylformamide (5ml) and treated with an excess of oxone (0.375g) and stirred at room temperature for 4h. The reaction was quenched by the addition of 1M aqueous sodium sulphite solution, then partitioned between dichloromethane and sodium bicarbonate solution. The solvent was removed *in vacuo* and the residue was purified by mass directed preparative HPLC (Method C). After
 20 evaporation of solvent the title compound was obtained as a yellow solid.

LC/MS R_t 2.30min, m/z 386 $[MH^+]$

The following were prepared in the same manner as Example 386:



Ex. No. (a)	R ³ SO ₂ -	R ²⁰ -	Starting Material	Thiol Reagent/ Source	LCMS MH ⁺	LCMS R _t (min)
384 (b)		H-	Intermediate 14	Isobutyl mercaptan/ Aldrich	414	2.56
387 HCl		H-	Intermediate 37	1-butanethiol/ Aldrich	414	2.67
388 HCl		H-	Intermediate 37	Methyl-3- mercaptopropionate / Fluka	444	2.39
389 HCl		H-	Intermediate 37	Phenethyl mercaptan/ Aldrich	462	2.88
390 HCl		H-	Intermediate 37	2- furanylmethanethiol / Aldrich	438	2.46
391 HCl		H-	Intermediate 37	2,2,2- trifluoroethanethiol/ Aldrich	440	2.6


392 HCl		Me-	Intermediate 45	N-(2-mercaptoethyl) acetamide/ Aldrich	457	2.25
------------	---	-----	-----------------	---	-----	------

(a) Salt forms: HCl = hydrochloride

(b) Isolation method: Mass Directed preparative HPLC (Method A).

5

Example 393. 3-[(3-(Aminocarbonyl)-8-methyl-4-{[3-(methyloxy)phenyl]amino}-6-quinolinyl)sulfonyl]benzoic acid




10

Example 379 (0.1g) was dissolved in methanol (5ml) and 2M aqueous sodium hydroxide (1ml). The mixture was heated to 75°C for 4h before cooling and standing at ambient temperature for 18h. The solvent was removed *in vacuo* and the residue partitioned between ethyl acetate (100ml) and water (100ml). The layers were separated and the aqueous layer was washed with diethyl ether (50ml), acidified to pH4 (2M hydrochloric acid) and extracted with ethyl acetate (2 x 150ml). The combined ethyl acetate layers were washed with brine (100ml), dried over magnesium sulfate and concentrated *in vacuo* to yield the title compound as a beige solid (0.082g).

LC/MS R_t 2.82min m/z 492 $[MH^+]$.

20

Similarly prepared were the following:




Example Number	R ³ SO ₂ -	Starting Material	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
394		Example 354	(II)	478	2.67
395		Example 355	(II)	478	2.65

(b) Isolation Method: (II) Mass Directed HPLC Method A




5

The following were prepared from the intermediates shown in the table in a similar manner to the method by which Example 393 was prepared, via Example 379, from Intermediate 45.



10

Example Number (a)	R ¹ NH-	Starting Material	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
--------------------	--------------------	-------------------	----------------------	----------------------	---------------------------

396 HCl		Intermediate 44	(IV)	516	2.25
397 HCl		Intermediate 46	(IV)	487	2.9
398		Intermediate 36	(I)	504	2.83


(a) Salt forms: HCl = hydrochloride

(b) Isolation Method:

(I) filtered off and used crude.

- 5 (IV) Mass Directed HPLC Method C; it is thought that compounds isolated by this method are hydrochloride salts.

Example 399. 6-({3-[(Dimethylamino)carbonyl]phenyl)sulfonyl}-8-methyl-4-{[3-(methoxy)phenyl]amino}-3-quinolinecarboxamide hydrochloride




- 15 To a solution of Example 393 (0.082g) in *N,N*-dimethylformamide (3ml) was added *N,N*-diisopropylethylamine (0.12ml) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (0.071g). The mixture was stirred for 20min before adding a solution of dimethylamine in tetrahydrofuran (2M, 0.8ml, Aldrich). After a

further 1h more dimethylamine in tetrahydrofuran (2M, 0.8ml) was added. After 1h the reaction mixture was concentrated *in vacuo* and partitioned between ethyl acetate (200ml) and aqueous sodium bicarbonate (100ml). The organic layers were washed with brine (100ml), dried over magnesium sulfate and concentrated *in vacuo*. Purification by mass

5 directed HPLC (Method C) gave the title compound as a yellow solid (0.022g).

LC/MS R_t 2.61min m/z 519 $[MH^+]$.

Similarly prepared were the following:



10

Example Number (a)	R¹NH-	R²⁰-	R²⁹R³⁰N-	Starting Material	Amine Reagent/ Source	LCMS MH⁺	LCMS R _t (min)
400 HCl		H-	Me₂N-	Example 395	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	505	2.52
401 HCl		H-	H₂N-	Example 395	Ammonia solution 880/ Merck	477	2.42
402 HCl		H-	MeNH-	Example 395	methylamine 2M solution in tetrahydrofuran/ Aldrich	491	2.47

403 HCl		H-		Example 395	Pyrrolidine/ Lancaster	531	2.64
404 HCl		H-		Example 395	propylamine/ Aldrich	519	2.74
405 (b)		H-		Example 395	Isobutylamine/ Aldrich	533	2.76
406 HCl		H-		Example 395	Morpholine/ Lancaster	547	2.51
407 HCl		H-		Example 395	Ethanolamine/ Aldrich	521	2.38
408 HCl		Me-	Me ₂ N-	Example 397	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	514	2.71
409 HCl		Me-	Me ₂ N-	Example 396	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	543	2.19
410 HCl		Me-	H ₂ N-	Example 396	Ammonia solution 880/ Merck	515	2.13
411 HCl		Me-	MeNH-	Example 396	methylamine 2M solution in tetrahydrofuran/ Aldrich	529	2.17
412 HCl		Me-	H ₂ N-	Example 397	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	486	2.61


413 HCl		Me-	MeNH-	Example 397	methylamine 2M solution in tetrahydrofuran/ Aldrich	500	2.69
414 (c)		Me-	Me ₂ N-	Example 398	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	531	2.65

(a) Salt form: HCl = hydrochloride



(b) Example 405 was isolated by Mass Directed preparative HPLC (Method A).

(c) Example 414 was isolated by aqueous work up.

5 Similarly prepared from Example 394 were the following:



Example Number (a)	R ²⁹ R ³⁰ N-	Amine Reagent/ Source	Isolation method (b)	LCMS MH ⁺	LCMS R _t (min)
415 HCl	H ₂ N-	Ammonia solution 880/ Merck	(I)	477	2.45
416 HCl	MeHN-	methylamine 2M solution in tetrahydrofuran/ Aldrich	(I)	491	2.51
417 HCl		Pyrrolidine/ Lancaster	(I)	531	2.66

418 HCl		propylamine/ Aldrich	(I)	519	2.75
419 HCl	Me ₂ N-	dimethylamine 2M solution in tetrahydrofuran/ Aldrich	(I)	505	2.53
420		Isobutylamine/ Aldrich	(II)	533	2.8


(a) Salt form: HCl = hydrochloride

(b) Isolation Method:

- (I) Mass Directed preparative HPLC (Method C).
- (II) Mass Directed preparative HPLC (Method A).

5

Example 421. 4-([3-(Methyloxy)phenyl]amino)-6-(4-piperidynylsulfonyl)-3-quinolinecarboxamide




10

To a mixture containing Example 472 (1.3g) in anisole (9ml) was added a solution of 95% trifluoroacetic acid in water (45ml). The mixture was stirred for 2h at room temperature and was then concentrated *in vacuo*. The residue was co-evaporated with toluene (2 x 20ml) and triturated with diethyl ether to give a yellow solid. The solid was partitioned



15 between aqueous potassium carbonate (300ml) and chloroform (300ml) and the aqueous phase extracted with chloroform (3 x 200ml). The combined organic extracts were washed with water (100ml), dried and concentrated *in vacuo* to give the title compound as a yellow solid (1.1g).

LC/MS R_t 1.94 min, *m/z* 441 [MH⁺]

Similarly prepared were the following:



Ex. No. (a)	R ¹ NH-	R ²⁰ -	R ³ SO ₂ -	Starting Material	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
422 TFA		Me-		Example 353	(I)	1.98	4.27
423		Me-		Example 352	(II)	4.67	1.97
470 TFA		Me-		Example 378	(III)	4.50	2.03
474 TFA		Me-		Example 376	(I)	4.79	1.77
476		Me-		Intermediate 55	(II)	4.73	2.10
477		Me-		Intermediate 56	(II)	4.33	2.03
561		Me-		Example 602	(IV)	4.03	1.93

567 HCl		Me-		Example 568	(V)	426	1.80
------------	---	-----	---	-------------	-----	-----	------

(a) Salt forms: TFA = trifluoroacetate

(b) Isolation Method:

(I) Filtered off directly from the reaction mixture; it is thought that compounds isolated by this method are trifluoroacetate salts.


(II) Aqueous workup of the crude reaction mixture without further purification; it is thought that compounds isolated by this method are free bases.

(III) Crude product was triturated to give the desired product and no further purification was carried out; it is thought that compounds isolated by this method are trifluoroacetate salts.

(IV) Product isolated by SCX ion exchange to give the free base

(V) Reaction mixture evaporated to dryness; it is assumed that this method gave the hydrochloride salt.

Example 424. 4-{[3-(Methoxy)phenyl]amino}-6-{[1-(phenylcarbonyl)-4-piperidyl]sulfonyl}-3-quinolinecarboxamide




To a mixture containing Example 421 (0.050g) and triethylamine (0.025ml) in dioxane (2ml) was added benzoyl chloride (0.020ml). The mixture was stirred under nitrogen for 18h at room temperature and was then diluted with methanol (5ml). The solution was applied to an aminopropyl cartridge and eluted with methanol. The eluent was evaporated and the residual gum purified by chromatography on SPE eluting with a

gradient of methanol in chloroform (0% to 10%) to give the title compound as a yellow solid (0.043g).

LC/MS R_t 2.63 min, m/z 545 $[MH^+]$

5 Similarly prepared were the following:



Ex. No. (a)	R^1NH-	$R^{20}-$	R^X-	Starting Material	Electrophile/ Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
425		H-		Example 421	Methyl chloroformate/ Aldrich	(II)	499	2.48
426		H-		Example 421	Acetyl chloride/ Aldrich	(II)	483	2.27
427		H-		Example 421	Methane sulphonyl chloride/ Aldrich	(V)	519	2.43
428		H-		Example 421	3-methylbutanoyl chloride/ Aldrich	(III)	525	2.61
429		H-		Example 421	Cyclopropane carbonyl chloride/ Aldrich	(III)	509	2.42

430		H-		Example 421	2-furancarboxyl chloride/ Aldrich	(III)	535	2.5
431		H-		Example 421	5-methyl-3-isoxazolecarboxyl chloride/ Maybridge	(III)	550	2.55
432		H-		Example 421	Benzene sulphonyl chloride/ Aldrich	(IV)	581	2.9
433		H-		Example 421	3,5-dimethyl-4-isoxazolesulfonyl chloride/ Avocado	(IV)	600	2.88
434		H-		Example 421	2-(acetylamino)-4-methyl-1,3-thiazole-5-sulfonyl chloride/ Aldrich	(IV)	659	2.69
435		H-		Example 421	1-butanethiol chloride/ Aldrich	(IV)	561	2.75
436		H-		Example 421	1-methylimidazole 4-sulphonyl chloride/ Maybridge	(IV)	585	2.35
437		H-		Example 421	Isoxazole -5-carboxyl chloride/ Lancaster	(III)	536	2.37
438		H-		Example 421	2-furancarboxyl chloride / Maybridge	(III)	535	2.42
439		H-		Example 421	Isobutyryl chloride/ Aldrich	(III)	511	2.42

440		H-		Example 421	Propionyl chloride/ Aldrich	(III)	497	2.31
441		H-		Example 421	1-pyrrolidinecarbonyl chloride/ Lancaster	(III)	538	2.44
442		Me-		Intermedi ate 54	2-furancarboxyl chloride / Aldrich	(I)	549	2.65
443		Me-		Example 470	2-furancarboxyl chloride / Aldrich	(I)	544	2.69
444		Me-		Example 470	Cyclopropane carbonyl chloride / Aldrich	(I)	518	2.63
445		Me-		Intermedi ate 54	Cyclopropane carbonyl chloride / Aldrich	(I)	523	2.57
446		Me-		Example 423	2-furancarboxyl chloride / Aldrich	(I)	561	2.66
447		Me-		Example 423	Cyclopropane carbonyl chloride / Aldrich	(I)	535	2.58
448 HCl		Me-		Example 423	Methyl chloroformate/ Aldrich	(VI)	525	2.57

(a) Salt forms: HCl = hydrochloride

(b) Isolation Method:

(I) Purified by chromatography on an SPE column.

(II) Aqueous workup of the crude reaction mixture without further purification.


(III) Purified using an SPE cartridge (aminopropyl solid phase) followed by chromatography using a silica SPE column.

(IV) Purified using an SPE cartridge (aminopropyl solid phase) followed by trituration.

(V) Aqueous workup of the crude reaction followed by trituration of the crude product.

5 (VI) Aqueous workup of the crude reaction followed by addition of dilute HCl in dioxane and evaporation; it is thought that compounds isolated by this method are hydrochloride salts.


10 **Example 449.** 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[(1-methyl-4-piperidinyl)sulfonyl]-3-quinolinecarboxamide hydrochloride



15 To a mixture containing Example 423 (0.050g) and triethylamine (0.025ml) in *N,N*-dimethylformamide (2ml) was added methyl iodide (0.0075ml). The mixture was stirred under nitrogen for 18h at room temperature and was concentrated by blowing down under nitrogen. Purification by chromatography on silica gel, eluting with dichloromethane/methanol (95:5), gave a white solid which was dissolved in dioxane (10ml) and treated with 4M hydrogen chloride in 1,4 dioxane (0.100ml). After evaporation
20 by blowing down under nitrogen the title compound was obtained as a yellow solid (0.028g).

LC/MS R_t 1.99 min, m/z 481 $[MH^+]$


25 **Example 450.** 6-[(1-Acetyl-4-piperidinyl)sulfonyl]-4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-3-quinolinecarboxamide hydrochloride



To a mixture containing Example 423 (0.050g) in pyridine (2ml) was added acetic anhydride (0.011ml). The mixture was stirred under nitrogen for 18h at room temperature, partitioned between chloroform (100ml) and 10% sodium carbonate solution (100ml), the layers separated by hydrophobic frit and the organic layer treated with 4M hydrogen chloride in 1,4-dioxane (0.100ml). After evaporation by blowing down under nitrogen the title compound was obtained as a pale yellow solid (0.021g).

LC/MS R_t 2.32 min, m/z 509 $[MH^+]$

10 Similarly prepared were the following:



Ex. No (a)	R^1NH-	$R^{20}-$	R^3SO_2-	Starting Material	Electrophile/Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
451		Me-		Example 476	Acetic anhydride/ Aldrich	(II)	515	2.31
452 HCl		Me-		Example 422	Acetyl chloride/ Aldrich	(V)	469	2.14

453 HCl		Me-		Example 477	Acetic anhydride/ Aldrich	(I)	475	2.15
454 HCl		Me-		Example 477	Methane- sulphonyl chloride/ Aldrich	(IV)	511	2.23
455 HCl		Me-		Example 477	Methyl chloroformate / Aldrich	(IV)	491	2.29
456		Me-		Example 422	Methyl chloroformate / Aldrich	(VI)	485	2.36
457		Me-		Example 422	Methane- sulphonyl chloride/ Aldrich	(VI)	505	2.31
531		Me-		Example 567	Acetic anhydride/ Aldrich	(III)	468	2.00

(a) Salt forms: HCl = hydrochloride.

(c) Isolation Method:

(I) As for Example 450

5 (II) Mass Directed HPLC (Method A)


(III) Purified by silica SPE eluting with ethyl acetate/ methanol

(IV) Mass Directed preparative HPLC (Method C).

(V) Aqueous work-up followed by addition of 4M HCl in 1,4-dioxane to a chloroform solution of the free base to give the hydrochloride salt.

10 (VI) Aminopropyl SPE column.


Example 473: 4-{[3-(Methyloxy)phenyl]amino}-6-({2-[2-(methylpropanoyl)amino]ethyl}sulfonyl)-3-quinolinecarboxamide



A solution of Example 364 (0.052g) in anisole (1ml) was treated with a solution of 95% trifluoroacetic acid in water (5ml). The mixture was stirred for 3h at room temperature and was then concentrated *in vacuo*. The residue was triturated with ethyl acetate, and the resulting solid was collected by filtration, washed with ethyl acetate and ether and dried to give a yellow solid (0.031g). The solid was treated with dioxane (2ml) and the suspension treated with *N,N*-diisopropylethylamine (0.04ml) followed by isobutyryl chloride (0.015ml, Aldrich) and the resulting solution stirred at room temperature under nitrogen for 2h. The solution was diluted with methanol (5ml) and applied to an aminopropyl SPE cartridge. Elution with methanol gave a gum after evaporation of solvent. The gum was purified by chromatography on silica gel eluting with a gradient of 0% to 6% methanol in chloroform to give the title compound as a yellow solid (0.017g).

LC/MS R_t 2.22 min, m/z 471 $[MH^+]$


Example 458. 6- $\{[1-(1H$ -imidazol-4-ylcarbonyl)-4-piperidiny]sulfonyl}-8-methyl-4- $\{[3-(methoxy)phenyl]amino$ }-3-quinolinecarboxamide



To a solution containing Intermediate 54 (0.041g) in *N,N*-dimethylformamide (3ml) were added imidazole-4-carboxylic acid (0.012g, Aldrich), (1*H*-1,2,3-benzotriazol-1-yloxy)(tri-1-pyrrolidinyl)phosphonium hexafluorophosphate (PyBop) (0.053g) and *N,N*-diisopropylethylamine (0.03ml). The solution was stirred under nitrogen at room temperature for 18h and was then concentrated *in vacuo*. The residual gum was purified using an aminopropyl SPE cartridge eluting with methanol followed by chromatography on silica gel (SPE cartridge), eluting with a gradient of 0% to 8% methanol in chloroform, to give the title compound as a yellow solid (0.031g).


LC/MS R_t 2.32 min, m/z 549 [MH^+]

Similarly prepared were the following:



Example Number (a)	R ^x -	Starting Material	Amine Reagent/ Source	LCMS MH ⁺	LCMS R _t (min)
459		Example 474	2-furoic acid/ Aldrich	573	2.15
460		Example 474	Cyclopropylmethanoic acid/ Aldrich	547	2.12


Example 461. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-{[1-(methylsulfonyl)-4-piperidinyl]sulfonyl}-3-quinolinecarboxamide hydrochloride



To a mixture containing Example 423 (0.050g) in 1,4-dioxane (2ml) was added methanesulphonyl chloride (0.009ml). The mixture was stirred under nitrogen for 18h at room temperature, partitioned between ethyl acetate (100ml) and 10% sodium bicarbonate solution (100ml), separated and dried. The solid obtained was dissolved in 1,4-dioxane and treated with 4M hydrogen chloride in 1,4-dioxane (0.100ml). After evaporation the title compound was obtained as a pale yellow solid (0.021g).

LC/MS R_t 2.5 min, m/z 545 $[MH^+]$


Example 462: 6-[[4-(Cyclopropylmethoxy)phenyl]sulfonyl]-4-[(3-methoxyphenyl)amino]quinoline-3-carboxamide hydrochloride



Tributyl phosphine (0.05ml) was added to a stirred mixture of Example 360 (0.052g), cyclopropylmethanol (0.028g) and di-*tert*-butylazodicarboxylate (0.06g) in tetrahydrofuran (1.5ml) at 20° under nitrogen, and stirring was continued at 20° for 3h. The solvent was concentrated and the residue purified by mass directed preparative HPLC (Method C) to give the title compound as a yellow solid (0.09g).

LC/MS R_t 3.4min, m/z 504 $[MH^+]$

Example 463: 6-[(4-Ethoxyphenyl)sulfonyl]-4-[(3-methoxyphenyl)amino]quinoline-3-carboxamide hydrochloride




.HCl

A stirred mixture of Example 360 (0.05g), iodoethane (0.35ml) and potassium carbonate (0.02g) in acetonitrile (1.5ml) was heated under reflux temperature for 1h. The solvent was evaporated to dryness. The resulting solid was partitioned between dichloromethane (2x15ml) and water (30ml). The extracts were dried (Na_2SO_4) and concentrated. The residual solid was purified by mass directed HPLC (Method C) to give the title compound as a pale yellow solid (0.033g).




LC/MS R_t 2.87 min, m/z 478 $[MH^+]$

The following examples were similarly prepared:




.HCl

Example Number	R^xO-	Starting Material	Alkylating Agent/	LCMS MH^+	LCMS R_t (min)
----------------	---------	-------------------	-------------------	-------------	------------------

(a)			Source		
464 HCl		Example 360	1-iodopropane/ Aldrich	492	3.07
465 HCl		Example 360	2-iodopropane/ Aldrich	492	3.00
466 HCl		Example 360	Iodocyclopenta- ne/ Aldrich	518	3.24

(a) Salt form: HCl = hydrochloride.


Example 467: 4-[[3-(3-Furyl)phenyl]amino]-6-[(4-methoxyphenyl)sulfonyl]quinoline-3-carboxamide hydrochloride



A stirred mixture of Example 254 (0.051g), 3-furanboronic acid (0.017g, Aldrich), tetrakis(triphenylphosphine) palladium(0) (0.05g) and 2M sodium carbonate solution (1ml) in dimethoxyethane (2ml) was heated at 100°C for 1h. The mixture was cooled and poured into 2M sodium carbonate solution and extracted into dichloromethane (2x15ml). The extracts were dried (Na₂SO₄) and concentrated. The residue was purified by mass directed HPLC (Method C) to give the title compound as a yellow solid (0.026g).

LC/MS R_t 2.93min, m/z 500 [MH⁺]

Similarly prepared were the following:




Ex. No.	R ¹ NH-	R ³ SO ₂ -	Starting Material	Boronic Acid/ Source	LCMS MH ⁺	LCMS R _t (min)
468 (a)			Example 366	[4-(methyloxy)phenyl] boronic acid/ Aldrich	540	3.18
469 (b)			Example 366	3-furanboronic acid/ Aldrich	500	3.02

(a) Example 468 was isolated as the free base by trituration with ether.

(b) Example 469 was isolated as the free base by chromatography on silica gel, eluting with ethyl acetate.

5

Example 475. 6-{[3-(Dimethylamino)-3-oxopropyl]sulfonyl}-4-{[4-fluoro-3-(methyloxy)phenyl]amino}-8-methyl-3-quinolinecarboxamide hydrochloride




10

A solution of Intermediate 53 (0.04g) in *N,N*-dimethylformamide (3ml) was treated with oxone (0.22g) and the resulting solution was stirred at room temperature overnight. The reaction was quenched by the addition on 1M sodium sulphite solution (1ml) and extracted into dichloromethane. The combined organic layers were dried using a

hydrophobic frit and evaporated *in vacuo*, and the product was dissolved in *N,N*-dimethylformamide (2ml) and treated with *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (0.016g). After 5 min, dimethylamine hydrochloride (0.065g) and *N,N*-diisopropylethylamine (0.015ml) in *N,N*-dimethylformamide (2ml) were added. The resulting solution was left standing at room temperature overnight. Chromatographic purification by SCX (IST Isolute™, 10g), eluting with methanol and 2M ammonia/methanol gave a yellow oil. Further purification by mass-directed HPLC (Method C) gave the title compound as a yellow solid (0.009g).

LC/MS R_t 2.34 min, m/z 489 [MH^+]


Example 540. 4-[(5-Chloro-3-pyridinyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide hydrochloride



To a solution of Intermediate 33 (0.050g) in *N,N*-dimethylformamide was added 5-chloro-3-pyridinamine (0.032g; Specs) and pyridine hydrochloride (0.029g) and the mixture heated at 90°C for 16h. The solvent was blown off under a stream of nitrogen at 45°C. The residue was triturated with acetonitrile and the resultant precipitate collected by filtration to give the title compound as a brown solid.

LC/MS R_t 2.25min m/z 391 [MH^+]

Similarly prepared were the following:



Ex. No. (a)	R ¹ NH-	R ³ SO ₂ -	Starting Material	Amine Reagent/ Source	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
539		MeSO ₂ -	Intermediate 33	5-Fluoro-3-pyridinamine/ Synchem OHG	(I)	375	2.2
541 HCl		MeSO ₂ -	Intermediate 33	6-Methyl-3-pyridinamine/ AsymChem	(IV)	371	1.79
543 HCl		MeSO ₂ -	Intermediate 33	4-Amino-2-benzofuran-1(3H)-one/ EP0529636A1	(II)	412	2.25
601		MeSO ₂ -	Intermediate 33	2,6-dimethyl-3-pyridylamine/ Lancaster	(III)	385	1.76

(a) Salt form: HCl = hydrochloride

5 (b) Isolation method:


(I) Trituration with acetonitrile followed by elution through an aminopropyl SPE cartridge with methanol.

(II) Reaction was performed at 80°C in acetonitrile and the product isolated by filtration of the reaction mixture.

10 (III) Mass Directed preparative HPLC (Method A) followed by chromatography on silica gel eluting with 3% methanol in dichloromethane.


(IV) Trituration with acetonitrile followed by isolation of the product by filtration.

Example 480. Ethyl 3-[[3-(aminocarbonyl)-4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-quinolinyl]sulfonyl]propanoate



- 5 To a solution of Intermediate 57 (0.82g) in *N,N*-dimethylformamide (25ml) was added oxone (4.5g). The mixture was stirred at room temperature for 2h before quenching with aqueous sodium sulphite solution and extracting with dichloromethane (2 x 25ml). The organic layers were combined, washed with water, dried using a hydrophobic frit,
- 10 concentrated *in vacuo* and purified by chromatography on silica gel, eluting with an ethyl acetate: cyclohexane gradient, to give the title compound as a yellow solid (0.12g). LC/MS R_t 2.61min m/z 484 [MH^+].

- 15 **Example 481.** 3-[[3-(Aminocarbonyl)-4-{4-fluoro-3-(methyloxy)phenyl}amino}-8-methyl-6-quinolinyl]sulfonyl]propanoic acid hydrochloride




To a solution of Intermediate 53 (0.8g) in *N,N*-dimethylformamide (10ml) was added oxone (4.6g). The mixture was stirred at room temperature for 48h before quenching with

aqueous sodium sulphite solution and extracting with dichloromethane (3 x 25ml). The aqueous layers were combined and applied to an Oasis cartridge, eluting with water and methanol. The methanol fractions were combined and concentrated *in vacuo*. The residue was applied to an SPE cartridge (Isolute, aminopropyl solid phase), eluting with methanol and 2M ammonia/methanol; evaporation of the methanol/ammonia fraction gave an orange oil. Further purification by mass directed preparative HPLC (Method C) gave the title compound as a yellow oil (0.003g).

LC/MS R_t 2.23min m/z 462 $[MH^+]$.


Similarly prepared were the following:



Example Number (a)	R ¹ NH-	Starting Material	LCMS MH ⁺	LCMS R _t (min)
549 HCl		Intermediate 59	432	2.38
550 HCl		Intermediate 60	415	1.92

(a) Salt forms: HCl = hydrochloride


Example 482. 4- $\{[4\text{-Fluoro-3-(methyloxy)phenyl}]\text{amino}\}$ -8-methyl-6- $\{[3\text{-(4-morpholinyl)-3-oxopropyl}]\text{sulfonyl}\}$ -3-quinolinecarboxamide hydrochloride







To a solution of Example 481 (0.035g) in *N,N*-dimethylformamide (2ml) was added *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (0.029g). After 5 min, morpholine (0.007ml, available from Aldrich) and *N,N*-diisopropylethylamine (0.026ml) were added. The resulting solution was stirred at room temperature overnight, and applied directly to an SCX cartridge (IST Isolute TM, 5g). Elution with methanol and 2M ammonia/methanol gave an orange residue, which was further purified by mass directed preparative HPLC (Method C) to give the title compound as a yellow solid (0.006g).

LC/MS R_t 2.37min m/z 531 $[MH^+]$.

Similarly prepared were the following:



Example Number (a)	R^3SO_2-	R^1NH-	Starting Material	Amine reagent / Source	Isolation Method (b)	LCMS MH^+	LCMS R_t (min)
483 HCl			Example 481	tetrahydro-2H-pyran-4-amine / Aldrich	(II)	545	2.22
484 HCl			Example 481	1-methyl piperazine / Aldrich	(II)	544	1.84


506 HCl			Example 550	1-methyl piperazine / Aldrich	(II)	497	1.75
507 HCl			Eample 549	1-methyl piperazine / Aldrich	(II)	514	1.98

(a) Salt form HCl = hydrochloride

(b) Isolation Method: (II) Mass Directed preparative HPLC (Method C).

5


Example 551. 6-[[3-(Dimethylamino)-3-oxopropyl]thio]-4-[(3-fluorophenyl)amino]-8-methyl-3-quinolinecarboxamide







To a solution of Intermediate 59 (0.04g) in *N,N*-dimethylformamide (1ml) was added *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (0.038g). After 5 min, dimethylamine hydrochloride (0.026g) and *N,N*-diisopropylethylamine (0.07ml) were added. The resulting solution was stirred at room temperature overnight, and applied directly to an SCX cartridge (IST Isolute™, 5g), eluting with methanol followed by 2M ammonia in methanol to give the title compound as an orange oil (0.038g).

LC/MS R_t 2.39min m/z 427 $[MH^+]$.


Similarly prepared were the following:



Example Number	R ²⁶ R ²⁷ N-	R ¹ NH-	Starting Material	Amine reagent / Source	LCMS MH ⁺	LCMS R _t (min)
552			Intermediate 59	tetrahydro-2H-pyran-4-amine / Aldrich	483	2.34
553			Intermediate 59	cyclopropyl(methyl)amine/ Karl Industries	453	2.56
554			Intermediate 59	Morpholine / Aldrich	469	2.36
555			Intermediate 59	Pyrrolidine / Aldrich	453	2.49
556			Intermediate 60	Dimethylamine / Aldrich	410	2.03
557			Intermediate 60	tetrahydro-2H-pyran-4-amine / Aldrich	466	2.01
558			Intermediate 60	cyclopropyl(methyl)amine/ Karl Industries	436	2.20

559			Intermediate 60	Morpholine / Aldrich	452	2.03
560			Intermediate 60	Pyrrolidine / Aldrich	436	2.13

Example 485. 6-[[3-(Dimethylamino)-3-oxopropylsulfonyl]-4-[(3-fluorophenyl)amino]-8-methyl-3-quinolinecarboxamide hydrochloride




5



















To a solution of the Example 551 (0.038g) in *N,N*-dimethylformamide (2ml) was added oxone (0.22g). The mixture was stirred at room temperature for 2h before quenching with aqueous sodium sulphite solution and extracting with dichloromethane. The organic layers were combined, dried by filtration through a hydrophobic frit and concentrated *in vacuo*. Purification by mass directed preparative HPLC (Method C) gave the title compound as a yellow solid (0.015g).





















10



















LC/MS R_t 2.31min m/z 459 $[MH^+]$.



















15 Similarly prepared were the following:















Example Number (a)	R ³ SO ₂ -	Starting Material	R ²⁰ -	R ¹ NH-	Isolation Method (b)	LCMS MH ⁺	LCMS R _t (min)
486 HCl		Example 552	Me-		(II)	515	2.39
487 HCl		Example 553	Me-		(II)	485	2.62
488 HCl		Example 554	Me-		(II)	501	2.38
489 HCl		Example 555	Me-		(II)	485	2.49
490 HCl		Example 556	Me-		(II)	442	1.98
491 HCl		Example 557	Me-		(II)	498	1.95
492 HCl		Example 558	Me-		(II)	468	2.1
493 HCl		Example 559	Me-		(II)	484	1.96
494 HCl		Example 560	Me-		(II)	468	2.04

525 HCOOH		Example 563	Me-		(I)	413	2.00
526		Example 564	Me-		(I)	430	2.50
527		Example 681	Me-		(III)	459	2.30
529		Example 565	Me-		(IV)	444	2.5
530 HCl		Example 566	Me-		(II)	427	2.00
532		Example 570	Me-		(V)	445	2.20
533 HCl		Example 571	Me-		(II)	442	2.3
534 HCOOH		Example 572	Me-		(I)	428	1.8
535 HCOOH		Example 573	Me-		(I)	467	2.6
538 HCl		Example 574	Me-		(II)	520	2.83

568		Example 569	Me-		(VI)	526	2.60
578 HCl		Example 635	Me-		(II)	385	2.13
579 HCl		Example 636	Me-		(II)	385	2.08
580 HCl		Example 637	Me-		(II)	399	2.22
581 HCl		Example 634	Me-		(II)	371	1.95
582 HCl		Example 638	Cl-		(II)	391	2.07
583 HCl		Example 662	Me-		(II)	388	2.33
584 HCl		Example 664	Me-		(II)	402	2.4
585 HCl		Example 665	Me-		(II)	416	2.54

666 HCl		Example 646	Cl-		(II)	425	2.52
667 HCl		Example 647	Cl-		(II)	439	2.73
668 HCl		Example 648	Cl-		(II)	439	2.56
669 HCl		Example 649	Cl-		(II)	453	2.66
670 HCl		Example 644	Me-		(II)	419	2.47
671 HCl		Example 650	Me-		(II)	389	2.28
672 HCl		Example 651	Me-		(II)	403	2.49
673 HCl		Example 652	Me-		(II)	403	2.35
674 HCl		Example 653	Me-		(II)	417	2.47

675 HCl		Example 655	Cl-		(II)	423	2.49
676 HCl		Example 656	Cl-		(II)	423	2.43
677 HCl		Example 657	Cl-		(II)	437	2.54
678 HCl		Example 642	Me-		(II)	405	2.38
679 HCl		Example 643	Me-		(II)	419	2.53
680 HCl		Example 645	Me-		(II)	433	2.67


(a) Salt form HCl = hydrochloride

HCOOH = formate

(b) Isolation Method:


- 5 (I) Mass Directed Preparative HPLC (Method A)
- (II) Mass Directed Preparative HPLC (Method C)
- (III) Aqueous work-up
- (IV) SCX ion exchange eluting with 2M ammonia in methanol
- (V) Trituration with methanol and collection of the product by filtration
- 10 (VI) Chromatography on silica gel eluting with methanol/ ethyl acetate mixtures

Example 495. 4-([4-Fluoro-3-(methyloxy)phenyl]amino)-8-methyl-6-([2-(2-oxo-1-pyrrolidiny)ethyl]sulfonyl)-3-quinolinecarboxamide hydrochloride



To a solution of Example 477 (0.03g) in 1,4-dioxan (5ml) was added ethyl 4-bromobutyrate (0.01ml, available from Aldrich). The mixture was heated at 120°C for 48h. The solvent was evaporated *in vacuo*. Purification by mass directed preparative HPLC (Method C) gave the title compound as a yellow solid (0.007g). LC/MS R_t 2.3min m/z 501 $[MH^+]$.


Similarly prepared were the following:



Example Number (a)	R ¹ NH-	Starting Material	LCMS MH ⁺	LCMS R _t (min)
514 HCl		Example 561	471	2.29

(a) Salt form HCl = hydrochloride

Example 518. 6-{[2-(Dimethylamino)ethyl]sulfonyl}-4-{[4-fluoro-3-(methoxy)phenyl]amino}-8-methyl-3-quinolinecarboxamide formate salt.



5

To a stirred mixture of Example 477 (0.05g) in dry *N,N*-dimethylformamide (1ml) was added methyl iodide (0.033g) and triethylamine (0.032ml), and the mixture was stirred under nitrogen for 18h. The mixture was applied directly to an SPE cartridge (1g) and eluted with 4% methanol in chloroform; the eluent was evaporated *in vacuo* and the residue purified using mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.003g).


10

LC/MS R_t 2.01min, m/z 461 [MH^+]

15

Example 519. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-6-{[2-(dimethylamino)ethyl]sulfonyl}-8-methyl-3-quinolinecarboxamide formate salt


Example 519 was prepared by a similar method to Example 518 from Example 422 to give the title compound as a yellow solid (0.005g)



20

LC/MS R_t 1.94min, m/z 455 [MH^+]

Example 521. 4-{[4-Fluoro-3-(methyloxy)phenyl]amino}-8-methyl-6-{[2-(methyloxy)ethyl]thio}-3-quinolinecarboxamide.



5

To a solution of Example 337 (0.05g) in dry *N,N*-dimethylformamide (2ml) under nitrogen was added sodium hydride (60% dispersion in mineral oil, 0.015g). The mixture was stirred at room temperature for 10min when methyl iodide (0.0078ml) was added; the mixture was stirred at room temperature for 18h and the solvent evaporated *in vacuo*. The residue was partitioned between chloroform and water, the layers separated by hydrophobic frit, and the organic layer evaporated. The crude product was purified using mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.025g).


LC/MS R_t 2.46min, m/z 416 $[MH^+]$

15

Similarly prepared from Example 528 was the following:


Example 571. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-{[2-(methyloxy)ethyl]thio}-3-quinolinecarboxamide

20



LC/MS R_t 2.40min, m/z 410 $[MH^+]$


Example 523. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-6-[(2-hydroxyethyl)sulfonyl]-8-methyl-3-quinolinecarboxamide.



To a solution of Example 528 (0.05g) in *N,N*-dimethylformamide (2ml) was added oxone (0.311g). The mixture was stirred at room temperature for 3h before quenching with aqueous sodium sulphite solution and extracting with ethyl acetate (50ml). The organic layer was dried (Na_2SO_4) and concentrated *in vacuo*, and the mixture purified by mass directed preparative HPLC (Method A) to give the title compound as a white solid (0.035g).

LC/MS R_t 2.1min m/z 428 $[MH^+]$.


Example 524. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[(2-methoxyethyl)sulfonyl]-3-quinolinecarboxamide



To a solution of Example 523 (0.018g) in dry *N,N*-dimethylformamide (1ml) under nitrogen was added sodium hydride (60% dispersion in mineral oil, 0.0017g). The mixture

was stirred at room temperature for 10min when methyl iodide (0.0026ml) was added, stirring was continued for 18h at room temperature and the solvent evaporated *in vacuo*. The residue was partitioned between ethyl acetate and water and the organic layer dried (MgSO₄) and evaporated. The crude product was purified using mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.0024g).
 LC/MS R_t 2.3min, *m/z* 442 [MH⁺]


Example 536. 6-({3-[(Dimethylamino)carbonyl]phenyl}sulfinyl)-8-methyl-4-{{3-(methoxy)phenyl}amino}-3-quinolinecarboxamide



To a mixture containing Example 544 (0.10g) in *N,N*-dimethylformamide (10ml) was added oxone (0.253g). The mixture was stirred under nitrogen for 3h at room temperature and was then quenched with a solution of sodium sulphite (0.25g) in water (10ml), diluted with water (30ml) and extracted with ethyl acetate (2 x 30ml). The combined organic extracts were evaporated to dryness and the residue purified by mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.028g).

LC/MS R_t 2.24 min, *m/z* 503 [MH⁺]


Example 537. 6-({3-[(Dimethylamino)carbonyl]phenyl}sulfonyl)-4-[(3-hydroxyphenyl)amino]-8-methyl-3-quinolinecarboxamide



A solution of borontribromide in dichloromethane (1.0M, 2.2ml) was added dropwise to an ice-cooled mixture containing Example 478 (0.35g) in dichloromethane (25ml) under nitrogen. The mixture was stirred at room temperature for 20h, and was then treated with a further portion of borontribromide in dichloromethane (1.0M, 2.2ml) and stirred for a further 5h. The mixture was quenched with methanol (10ml) and evaporated to dryness *in vacuo*. The residue was partitioned between ethyl acetate (30ml) and saturated aqueous sodium bicarbonate (30ml), the organic extract evaporated to dryness *in vacuo*, and the residue purified by mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.075g).

LC/MS R_t 2.46 min, m/z 505 $[MH^+]$

Example 575: 7-({3-[(Dimethylamino)carbonyl]phenyl}sulfinyl)-4-{{3-(methoxy)phenyl}amino}-3-quinolinecarboxamide




A mixture containing Intermediate 65 (0.15g), 10% palladium on activated carbon (0.04g) and triethylamine (5ml) in ethanol (25ml) and *N,N*-dimethylformamide (10ml) was hydrogenated at room temperature for 4h. The suspension was filtered through celite, the residue washed with ethanol / *N,N*-dimethylformamide (3:1, 50ml), and the filtrate

concentrated *in vacuo*. The residue was purified by mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.035g).

LC/MS R_t 2.32min, m/z 489 $[MH^+]$

5

Example 545: 7-({3-[(Dimethylamino)carbonyl]phenyl}sulfonyl)-4-{[3-(methyloxy)phenyl]amino}-3-quinolinecarboxamide



10


Oxone (0.22g) was added portionwise to a stirred solution of Example 575 (0.035g) in *N,N*-dimethylformamide (4ml). The mixture was stirred at room temperature under nitrogen for 24h, a further portion of oxone (0.17g) was added, and the mixture stirred for a further 5h. The reaction was quenched with a solution of sodium sulphite (1.2g) in water (15ml), diluted with water (10ml) and extracted with ethyl acetate (3 x 30ml). The combined organic extracts were dried over magnesium sulphate and concentrated *in vacuo* to give the title compound as a buff solid (0.035g).

15

LC/MS R_t 2.66min, m/z 505 $[MH^+]$

20

Example 576: 6-({5-[(Dimethylamino)carbonyl]-3-pyridinyl}thio)-8-methyl-4-{[3-(methyloxy)phenyl]amino}-3-quinolinecarboxamide, formate salt





25

A stirred mixture of Intermediate 45 (0.47g), Intermediate 69 (0.37g), copper iodide (0.06g), and potassium carbonate (0.47g) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (10ml) was heated at 100°C under nitrogen for 4h. The mixture was diluted with water (150ml) and extracted with ethyl acetate (3 x 200ml). The combined organic extracts were washed with water (2 x 200ml) and brine (200ml), and the organic layers dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified by mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.1g).

LC/MS R_t 2.35min, m/z 488 [MH^+]

Example 547. 6-({5-[(Dimethylamino)carbonyl]-3-pyridinyl}sulfinyl)-8-methyl-4-{[3-(methyloxy)phenyl]amino}-3-quinolinecarboxamide hydrochloride and

Example 546. 6-({5-[(Dimethylamino)carbonyl]-3-pyridinyl}sulfonyl)-8-methyl-4-{[3-(methyloxy)phenyl]amino}-3-quinolinecarboxamide hydrochloride




Oxone (1.2g) was added portionwise to a stirred solution of Example 576 (0.1g) in *N,N*-dimethylformamide (6ml). The solution was stirred at room temperature under nitrogen for 2h and then quenched with a solution of sodium sulphite (3g) in water (30ml). The mixture was diluted with water (25ml) and extracted with ethyl acetate (4 x 50ml) and the combined organic layers were washed with water (2 x 50ml) and brine (50ml), dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified using mass directed preparative HPLC (Method C) to give Example 546 as a yellow solid (0.010g) and Example 547 as a yellow solid (0.041g).

Example 546: LC/MS R_t 2.57min, m/z 520 [MH^+]

Example 547: LC/MS R_t 2.12min, m/z 504 [MH^+]


Example 586. 8-Methyl-4-[(3-methyl-5-isoxazolyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide hydrochloride



To a stirred suspension of sodium hydride (0.008g; 60% dispersion in mineral oil) in dry *N,N*-dimethylformamide (1ml) was added [(3-methyl-5-isoxazolyl)methyl]amine (available from Aldrich) (0.020g) and the mixture heated at 80°C for 30 min. A suspension of Intermediate 33 (0.020g) in dry *N,N*-dimethylformamide (0.5ml) was added and the mixture heated at 80°C for 3h. The mixture was quenched by the dropwise addition of ethanol (0.1ml). The mixture was loaded onto a 2g SCX cartridge, washed with methanol, and the product eluted with 10% '880' ammonia in methanol. The solvent was removed *in vacuo* and the residue purified by mass directed preparative HPLC (Method C) to give the title compound as a pale yellow solid (0.009g)

LC/MS R_t 2.23min, m/z 361 [MH^+]


Example 544. 6-({3-[(Dimethylamino)carbonyl]phenyl}thio)-8-methyl-4-[(3-(methoxy)phenyl)amino]-3-quinolinecarboxamide



A stirred mixture of Intermediate 45 (50g), Intermediate 28 (40g), and potassium carbonate (40g) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (500ml) was
 5 purged of air (by evacuation of the vessel and refilling with nitrogen three times) and left under nitrogen. Copper (I) iodide (5g) was added and the mixture was warmed at 90°C for 23h. The mixture was cooled to 20°C and poured into water (2.5L). The precipitated solid was filtered off, washed with water and sucked partially dry. The damp solid was dissolved in chloroform (4L) and washed with 1N sodium hydroxide solution (1L), followed
 10 by water (2 x 1L) and brine (1L). The organic phase was dried over sodium sulphate and the solvent evaporated to leave a sticky solid. The solid was crystallised from hot ethanol (650ml) to give the title compound as a solid (45.1g).

LC/MS R_t 2.60min m/z 487 $[MH^+]$.

Similarly prepared were the following:



15

Example Number (a)	R ¹ NH-	R ³ S-	Starting Material	Isolation method (b)	LCMS MH ⁺	LCMS R _t (min)
562			Intermediate 45	(I)	417	2.27

573 HCl			Intermediate 35	(II)	435	2.56
574			Intermediate 36	(III)	487	2.86

(a) Salt form: HCl = hydrochloride


(b) Isolation method: (I) Aqueous work-up followed by trituration with ether and filtration.

(II) Mass directed preparative HPLC (Method C).

(III) Trituration with ether and filtration.

5

Example 544. 6-({3-[(Dimethylamino)carbonyl]phenyl}thio)-8-methyl-4-{[3-(methyloxy)phenyl]amino} -3-quinolinecarboxamide (alternative synthesis)



10


Intermediate 45 (5.0g), Intermediate 28 (2.89g), copper iodide (0.506g) and potassium carbonate (2.94g) were added to 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU, 25ml) and the resulting stirred slurry was heated to 100°C under nitrogen. The mixture was stirred at 100°C for 7h, allowed to cooled to room temperature and stirred overnight. DMPU (20ml) and water (80ml) containing pyridine (0.43ml) were added and the slurry was heated to 100°C. The resulting solution was seeded with crystals of Example 544 and stirred for 1h at 100°C. The suspension was cooled gradually over 6h, allowing the product to crystallise. The product was isolated by filtration, washed with water (2x 50ml) and dried at 40°C *in vacuo* to give the title compound as a pale yellow solid (3.9g).

15

20

LC/MS R_t 2.58min m/z 487 $[MH^+]$.

Example 478. 6-({3-[(Dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-{{3-(methoxy)phenyl} amino}-3-quinolinecarboxamide (alternative synthesis)



5

To a solution of Example 544 (29g) in *N,N*-dimethylformamide (290ml) cooled in a water bath was added oxone (87g) in portions over 10min. The mixture was stirred for 2h, then poured into a cold (5°C) solution of sodium metabisulphite (45g) in water (2L). After stirring for 35min the mixture was extracted with chloroform (2L + 3 x 800ml). The combined chloroform extracts were washed with water (3 x 600ml), and the aqueous washes were extracted with chloroform (600ml). The combined organic phases were dried over sodium sulphate and the solvent evaporated to leave a solid which was dried *in vacuo* at 40°C for 3 days providing the title compound (27.8g).

10

LC/MS R_t 2.62min m/z 519 [MH^+].


15

The solid was crystallised from hot ethanol containing 20% water (5L) to give the title compound (20.2g).

LC/MS R_t 2.62min m/z 519 [MH^+].

20

Example 478. 6-({3-[(Dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-{{3-(methoxy)phenyl} amino}-3-quinolinecarboxamide (alternative procedure)




To a solution of Example 544 (3.5g) in glacial acetic acid (18ml) and water (3.5ml) was added oxone (5.76g) portionwise over 15 min. The mixture was stirred for 1.5h at 20°C and excess oxone quenched with a solution of sodium sulphite (0.545g) in water (3.5ml).

- 5 The mixture was diluted with glacial acetic acid (11ml) and water (21ml), heated to 90°, treated dropwise over 30 min with 2M aqueous sodium hydroxide (20ml), and cooled to 25°C over 30min. The resulting precipitate was collected by filtration, washed with water (25ml x3) and dried *in vacuo* to give the title compound as a pale yellow solid (3.0g).
LC/MS R_t 2.54min m/z 519 $[MH^+]$.

10

Example 588: 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfinyl)-3-quinolinecarboxamide formate salt.




15

To a suspension of Example 577 (0.04g) in methanol (10ml) was added sodium periodate (0.023g) in water (0.2ml). The mixture was stirred at room temperature for 4 days and the solvents evaporated *in vacuo*. The residue was purified using mass directed preparative HPLC (Method A) to give the title compound as a yellow solid (0.017g).

- 20 LC/MS R_t 2.0min, m/z 382 $[MH^+]$


Example 307: 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide hydrochloride (alternative synthesis).



- 5 To a solution of Example 577 (0.04g) in *N,N*-dimethylformamide (1ml) was added oxone (0.337g). The mixture was stirred at room temperature for 18h, and quenched by addition of 10% sodium sulphite solution (15ml). The mixture was extracted with ethyl acetate (50ml), and the organic layer dried (Na_2SO_4) and evaporated *in vacuo*. The residue was purified using mass directed preparative HPLC (Method C) to give the title compound as a yellow solid (0.018g).
- 10

LC/MS R_t 2.2min, m/z 398 [MH^+]

- 15 **Example 688.** 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-7-(methylthio)-3-quinolinecarboxamide.




- 20 A stirred mixture of Intermediate 104 (0.50g), sodium methanethiolate (0.35g), potassium carbonate (0.43g) and copper (I) iodide (0.025g) in dry *N,N*-dimethylformamide (3ml) was heated at 100° under nitrogen for 18h. The mixture was cooled, poured into water (50ml) and stirred for 15min. The solid material was filtered off, dried *in vacuo* at 80° for 2h, and boiled in ethanol:water 15:1 (50ml) for 30min. The insoluble material was filtered off, and

the filtrate evaporated to dryness to give the title compound as a pale yellow solid (0.163g).

LC/MS R_t 2.40min m/z 366 $[MH^+]$

5

Example 548. 4-(2,3-Dihydro-1-benzofuran-4-ylamino)-8-methyl-7-(methylsulfonyl)-3-quinolinecarboxamide hydrochloride.




- 10 Example 548 was prepared from Example 688 by a similar method to Example 129, but without the addition of anisole to the reaction mixture, using 10:1 *N,N*-dimethylformamide: water as solvent, and purifying by mass directed preparative HPLC (method C).

LC/MS R_t 2.50min m/z 398 $[MH^+]$

CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein:

R¹ is

C₁₋₆ alkyl;

C₃₋₇cycloalkyl or C₃₋₇cycloalkyl(C₁₋₄alkyl)- wherein the C₃₋₇cycloalkyl is optionally substituted by one or more substituents selected from =O and OH;

C₄₋₇cycloalkyl fused to an aryl ring;

Aryl or aryl(C₁₋₆alkyl)- wherein the aryl is optionally substituted by one or more substituents selected from C₁₋₆alkyl, C₁₋₆alkylCONR⁶-, C₁₋₆alkylCO-, halogen, -CF₃, -(CH₂)_mOH, -OCF₃, C₁₋₆alkoxy-, C₁₋₆alkoxy(C₁₋₄alkyl)-, C₁₋₆alkoxyC₂₋₆alkoxy-, C₁₋₆alkoxycarbonyl, -CN, R⁴R⁵NCO, R⁷R⁸N-, R⁹R¹⁰NCONR¹¹-, HO(CH₂)₂₋₆O-, R¹²R¹³NSO₂(CH₂)_m-, (4-morpholinyl)C₂₋₆alkoxy, -NR¹⁴SO₂C₁₋₆alkyl, aryloxy, heteroaryl (optionally substituted by C₁₋₆alkyl), CO₂H, R²¹R²²N(C₁₋₄alkyl)-, C₁₋₆alkoxyCONR²³(CH₂)_m-, aryl(optionally substituted by C₁₋₆alkyl);

Aryl fused to a C₄₋₇cycloalkyl ring, wherein the cycloalkyl ring is optionally substituted by one or more =O;

5 Aryl fused to a heterocyclyl ring, wherein the heterocyclyl ring is optionally substituted by one or more substituents selected from =O, -COC₁₋₄alkyl, C₁₋₄alkyl;

Heteroaryl or heteroaryl(C₁₋₆alkyl)- wherein the heteroaryl is optionally substituted by one or more substituents selected from: C₁₋₆alkyl, aryl(C₁₋₄alkyl), C₁₋₆alkoxy, halogen, C₁₋₆alkoxyCO; or


10

Heterocyclyl optionally fused to an aryl or heteroaryl ring.

R² is hydrogen or C₁₋₆alkyl;

15

R³⁴ is hydrogen or a group of formula:





wherein R³ is

20

C₁₋₆alkyl optionally substituted by one or more substituents selected from -OH, -NR¹⁶COR¹⁵, -NR¹⁷R¹⁸, -CO₂R²⁴, C₁₋₆alkoxyCONR²⁵-, -CONR²⁶R²⁷, C₁₋₆alkoxy-, C₁₋₆alkylSO₂NR³³-, or a group having one of the following formulae;

25



C₃₋₇cycloalkyl;

Aryl or aryl(C₁₋₆alkyl)- wherein the aryl is optionally substituted by one or more substituents selected from C₁₋₆alkyl-, halogen-, C₁₋₆alkoxy-, -CO₂R²⁸, -CH₂CO₂H, -OH, aryl(optionally substituted by a C₁₋₆ alkoxy group), heteroaryl, -CONR²⁹R³⁰, C₃₋₇cycloalkoxy, C₃₋₇cycloalkyl(C₁₋₆alkoxy)-, -CF₃;

5

Heteroaryl or heteroaryl(C₁₋₆alkyl)- wherein the heteroaryl is optionally substituted by one or more C₁₋₆alkyl or -CONR²⁹R³⁰ groups; or

10

Heterocyclyl which is optionally substituted by one of more substituents selected from C₁₋₆alkyl-, C₁₋₆alkylCO-, C₃₋₇cycloalkylCO-, heteroarylCO- (optionally substituted by one or more C₁₋₄alkyl- groups), C₁₋₆alkoxyCO-, arylCO-, R³¹R³²NCO-, C₁₋₆alkylSO₂-, arylSO₂-, -heteroarylSO₂ (optionally substituted by one or more C₁₋₄alkyl or C₁₋₄alkylCONH- groups)

The heterocyclyl is linked to the S(=O)_n moiety through a carbon atom.


15

m is 0-6

n is 0, 1 or 2;

20

R¹⁹ is hydrogen, C₁₋₆alkyl or a group of formula:



R²⁰ is hydrogen, C₁₋₆alkyl, halogen or C₁₋₆alkoxy.

25

R⁴⁻¹⁸, R²¹⁻²⁵, R²⁸ and R³¹⁻³³ all independently represent H, C₁₋₆ alkyl;

R²⁶ and R²⁷ independently represent H, C₁₋₆ alkyl, C₃₋₇cycloalkyl or heterocyclyl

30

R²⁹ and R³⁰ independently represent H, C₁₋₆alkyl optionally substituted by OH;

R⁷ and R⁸ together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

5 R⁹ and R¹⁰ together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

R¹⁷ and R¹⁸ together with the nitrogen atom to which they are attached may form a heterocyclyl ring such as morpholine;

10 R²¹ and R²² together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

R²⁶ and R²⁷ together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

15 R²⁹ and R³⁰ together with the nitrogen atom to which they are attached may form a heterocyclyl ring such as morpholine;

20 R³¹ and R³² together with the nitrogen atom to which they are attached may form a heterocyclyl ring;

with the proviso that R³⁴ and R¹⁹ cannot both represent R³S(=O)_n.

2. A compound according to claim 1 wherein R¹ is selected from




25 Aryl optionally substituted by one or more substituents selected from C₁₋₆alkyl, halogen, C₁₋₆alkoxy-, -CN, -(CH₂)_m(OH), C₁₋₆alkylCO-;

Aryl fused to a heterocyclyl ring,

30 Heteroaryl optionally substituted by one or more substituents selected from: C₁₋₆alkyl, halogen, or C₁₋₆alkoxy groups.

3. A compound according to claim 1 or claim 2 wherein R¹ is selected from benzothiazolyl, benzimidazolyl, indazolyl, pyridyl and pyrazolyl.

4. A compound according to any one of the preceding claims wherein R¹ is selected from



5. A compound according to any one of the preceding claims wherein R¹ is selected from

Aryl optionally substituted by one or more substituents selected from: methyl, fluorine, chlorine, -CN, -OMe, -OH, COMe

Heteroaryl optionally substituted by one or more methyl, ethyl, fluorine, chlorine or methoxy groups.

6. A compound according to any one of the preceding claims wherein R¹ is selected from 3-(methyloxy)phenyl, 3-methylphenyl, 3-cyanophenyl, 3-fluorophenyl, 3-chlorophenyl, 4-fluoro-3-(methyloxy)phenyl, 3-acetylphenyl, 4-hydroxy-3-(methyloxy)phenyl, 2-fluoro-3-chlorophenyl, 2,3-difluorophenyl, 3,5-difluorophenyl, 1-

methyl-1H-benzimidazolyl-6-yl, 1-methyl-1H-indazol-6-yl, 5-(methyloxy)-3-pyridinyl, 3-pyridinyl, 1-ethyl-1H-pyrazol-5-yl, 5-methyl-3-pyridinyl, 1,3-benzothiazol-6-yl, 5-fluoro-3-pyridinyl, 5-chloro-3-pyridinyl.

- 5 7. A compound according to any one of the preceding claims wherein R² is hydrogen.
8. A compound according to any one of the preceding claims wherein R³ is selected from:
- 10 C₁₋₆ alkyl which is optionally substituted by one or more substituents selected from -NR¹⁶COR¹⁵; -CONR²⁶R²⁷,
- C₃₋₇cycloalkyl;
- 15 Aryl optionally substituted by one or more substituents selected from C₁₋₆alkyl-, C₁₋₆alkoxy-, CONR²⁹R³⁰;
- Heterocyclyl which is optionally substituted by one or more substituents selected from C₁₋₆alkylCO-, C₃₋₇cycloalkylCO-, heteroarylCO-;
- 20 9. A compound according to any one of the preceding claims wherein R³ is selected from methyl, ethyl, n-propyl, *tert*-butyl, isopropyl, MeCONH(CH₂)₂-, Me₂NCO(CH₂)₂-;
- Cyclopentyl;
- 25 Aryl optionally substituted by one or more methoxy, methyl, -CONMe₂ groups;
- Heterocyclyl which is optionally substituted by one of more substituents selected from MeCO-, cyclopropylCO, 2-furylCO-.
- 30 10. A compound according to any one of the preceding claims wherein R³ is 4-(methyloxy)phenyl, phenyl, 3-[(dimethylamino)carbonyl]phenyl, 4-methylphenyl, 3,4-bis(methyloxy)phenyl, 3,4,5-tris(methyloxy)phenyl, 3-(ethyloxy)phenyl.

11. A compound according to any one of the preceding claims wherein R³ is a piperidiny group substituted by one or more substituents selected from MeCO-, cyclopropylCO-, 2-furylCO-.


5

12. A compound according to any one of the preceding claims wherein R³ is 1-acetyl-4-piperidiny, 1-(2-furanylcarbonyl)-4-piperidiny, or 1-(cyclopropylcarbonyl)-4-piperidiny.

10 13. A compound according to any one of the preceding claims wherein R²⁰ is hydrogen, methyl, ethyl, fluorine or chlorine.

14. A compound according to any one of the preceding claims wherein n is 2.

15 15. A compound according to any one of the preceding claims wherein R³⁴ represents a group of formula:



20 16. A compound according to claim 1 which is

4-[(3-methylphenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-[[3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

25 4-[[4-fluoro-3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-[(3-chlorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-(1,3-benzothiazol-6-ylamino)-6-(phenylsulfonyl)-3-quinolinecarboxamide,

4-[(1-methyl-1H-benzimidazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

30 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(phenylsulfonyl)-3-quinolinecarboxamide,

4-[[3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,

- 6-(cyclopentylsulfonyl)-4-[(3-fluorophenyl)amino]-3-quinolinecarboxamide,
4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide,
6-[(1,1-dimethylethyl)sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-
5 quinolinecarboxamide,
6-[[2-(acetylamino)ethyl]sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-
quinolinecarboxamide.
6-[(1,1-dimethylethyl)thio]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,
6-[[2-(acetylamino)ethyl]thio]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,
10 4-[(1-methyl-1H-indazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,
4-[[4-hydroxy-3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,
4-[(3-acetylphenyl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,
8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide,
4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-6-(phenylsulfonyl)-3-
15 quinolinecarboxamide,
7-methyl-4-[[3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,
8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide,
4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
20 quinolinecarboxamide,
4-[(3-acetylphenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide
8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide
25 4-(2,3-dihydro-1,4-benzodioxin-5-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-
3-quinolinecarboxamide
4-[(3-chlorophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide
4-[(3-cyanophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
30 quinolinecarboxamide
4-(1,3-benzothiazol-6-ylamino)-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-
quinolinecarboxamide

- 4-[(3-fluorophenyl)amino]-8-methyl-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide
- 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide
- 5 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide
- 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[(4-methylphenyl)sulfonyl]-3-quinolinecarboxamide
- 4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-6-(methysulfonyl)-3-quinolinecarboxamide
- 10 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(phenylsulfonyl)-3-quinolinecarboxamide
- 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methysulfonyl)-3-quinolinecarboxamide
- 15 8-methyl-6-(methysulfonyl)-4-(3-pyridinylamino)-3-quinolinecarboxamide
- 8-methyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 4-[(3-fluorophenyl)amino]-8-methyl-6-(methysulfonyl)-3-quinolinecarboxamide
- 4-[(3-cyanophenyl)amino]-8-methyl-6-(methysulfonyl)-3-quinolinecarboxamide
- 20 4-[(1-ethyl-1H-pyrazol-5-yl)amino]-8-methyl-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-methyl-4-[[5-(methyloxy)-3-pyridinyl]amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-methyl-4-[(5-methyl-3-pyridinyl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 25 8-chloro-4-[(3-methylphenyl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-chloro-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-chloro-4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-(methysulfonyl)-3-quinolinecarboxamide
- 30 8-chloro-4-[(3-cyanophenyl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-chloro-4-[(3-fluorophenyl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide
- 8-chloro-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(methysulfonyl)-3-quinolinecarboxamide

- methyl 3-[(3-(aminocarbonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-quinoliny]sulfonyl]benzoate
- 6-[[3,4-bis(methyloxy)phenyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 5 8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[3,4,5-tris(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide hydrochloride
- 6-[[3,4-bis(methyloxy)phenyl]sulfonyl]-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 6-[[3-(ethyloxy)phenyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 10 6-[[2-(acetylamino)ethyl]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,
- 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,
- 15 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,
- 4-[(3-cyanophenyl)amino]-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide,
- 6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[(1-methyl-1H-benzimidazol-6-yl)amino]-3-quinolinecarboxamide
- 20 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide
- 6-[(1-acetyl-4-piperidiny]sulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 25 6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 4-[(3-cyanophenyl)amino]-6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-3-quinolinecarboxamide
- 6-[[1-(cyclopropylcarbonyl)-4-piperidiny]sulfonyl]-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide
- 30 4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-[[1-(2-furany]carbonyl)-4-piperidiny]sulfonyl]-8-methyl-3-quinolinecarboxamide

- 6-[[1-(cyclopropylcarbonyl)-4-piperidinyl]sulfonyl]-4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-3-quinolinecarboxamide
- 6-[[1-acetyl-4-piperidinyl]sulfonyl]-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-3-quinolinecarboxamide
- 5 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-({2- [(methylsulfonyl) amino]ethyl}sulfonyl)-3-quinolinecarboxamide
- 6-[[1-(2-furanylcarbonyl)-4-piperidinyl]sulfonyl]-8-methyl-4-[(1-methyl-1*H*-benzimidazol-6-yl)amino]-3-quinolinecarboxamide
- 10 6-[[3-(dimethylamino)-3-oxopropyl]sulfonyl]-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-8-methyl-3-quinolinecarboxamide
- 4-[(2,3-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-[(3-chloro-2-fluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-[(3,5-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 15 4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 6-({5-[(dimethylamino)carbonyl]-3-pyridinyl}sulfonyl)-8-methyl-4-[[3- (methyloxy) phenyl]amino]-3-quinolinecarboxamide hydrochloride
- 8-methyl-6-[(1-methylethyl)sulfonyl]-4-(3-pyridinylamino)-3-quinolinecarboxamide
- 20 6-[(1,1-dimethylethyl)sulfonyl]-8-methyl-4-(3-pyridinylamino)-3-quinolinecarboxamide
- 4-[(1-ethyl-1*H*-pyrazol-5-yl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide
- 6-[(1,1-dimethylethyl)sulfonyl]-4-[(1-ethyl-1*H*-pyrazol-5-yl)amino]-8-methyl-3-quinolinecarboxamide
- 25 4-(3,4-dihydro-2*H*-chromen-5-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-3-quinolinecarboxamide
- 30 8-ethyl-4-[[4-fluoro-3-(methyloxy)phenyl]amino]-6-(methylsulfonyl)-3-quinolinecarboxamide
- 8-ethyl-4-[(3-fluorophenyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide
- 4-[(3-cyanophenyl)amino]-8-ethyl-6-(methylsulfonyl)-3-quinolinecarboxamide

- 8-ethyl-4-[(1-methyl-1H-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide
 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-ethyl-6-(methylsulfonyl)-3-quinolinecarboxamide
 5 8-ethyl-6-(methylsulfonyl)-4-(3-pyridinylamino)-3-quinolinecarboxamide
 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-fluoro-6-(methylsulfonyl)-3-quinolinecarboxamide
 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-(ethylsulfonyl)-3-quinolinecarboxamide
 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-(propylsulfonyl)-3-quinolinecarboxamide
 10 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide
 8-chloro-4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-3-quinolinecarboxamide
 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide
 15 6-(ethylsulfonyl)-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide
 6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide
 8-chloro-4-[(5-fluoro-3-pyridinyl)amino]-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide
 20 8-chloro-6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-3-quinolinecarboxamide
 4-[(5-chloro-3-pyridinyl)amino]-6-(ethylsulfonyl)-8-methyl-3-quinolinecarboxamide
 4-[(5-chloro-3-pyridinyl)amino]-8-methyl-6-(propylsulfonyl)-3-quinolinecarboxamide
 25 4-[(5-chloro-3-pyridinyl)amino]-6-[(1,1-dimethylethyl)sulfonyl]-8-methyl-3-quinolinecarboxamide

or a pharmaceutically acceptable salt thereof.

- 30 17. A compound according to Claim 1 which is
 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

8-methyl-4-[(1-methyl-1*H*-indazol-6-yl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide,

8-methyl-4-[(5-methyl-3-pyridinyl)amino]-6-(methylsulfonyl)-3-quinolinecarboxamide

4-[(3,5-difluorophenyl)amino]-8-methyl-6-(methylsulfonyl)-3-quinolinecarboxamide

4-[(1-ethyl-1*H*-pyrazol-5-yl)amino]-8-methyl-6-[(1-methylethyl)sulfonyl]-3-quinolinecarboxamide

6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-8-methyl-3-quinolinecarboxamide

8-chloro-6-[(1,1-dimethylethyl)sulfonyl]-4-[(5-fluoro-3-pyridinyl)amino]-3-quinolinecarboxamide

or a pharmaceutically acceptable salt thereof.

18. A compound according to Claim 1 which is

8-methyl-4-[[3-(methyloxy)phenyl]amino]-6-[[4-(methyloxy)phenyl]sulfonyl]-3-quinolinecarboxamide,

6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide hydrochloride

6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide,

4-[(3-cyanophenyl)amino]-6-({3-[(dimethylamino)carbonyl]phenyl}sulfonyl)-8-methyl-3-quinolinecarboxamide,

4-(2,3-dihydro-1-benzofuran-4-ylamino)-6-[[1-(2-furanylcarbonyl)-4-piperidinyl]sulfonyl]-8-methyl-3-quinolinecarboxamide

5 4-(2,3-dihydro-1-benzofuran-4-ylamino)-8-methyl-6-({2-[(methylsulfonyl)amino]ethyl}sulfonyl)-3-quinolinecarboxamide

6-({5-[(dimethylamino)carbonyl]-3-pyridinyl}sulfonyl)-8-methyl-4-[[3-(methyloxy)phenyl]amino]-3-quinolinecarboxamide hydrochloride

10

or a pharmaceutically acceptable salt thereof.

19. A method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective PDE4 inhibitor is indicated, which comprises
15 administration of a therapeutically effective amount of a compound of formula (I) according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof.

20. A compound of formula (I) according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof for use in medical therapy.

20

21. A compound according to claim 20 for use in the treatment of inflammatory and/or allergic diseases.

22. A pharmaceutical formulation comprising a compound of formula (I) according to any
25 of claims 1 to 18 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.


23. A combination comprising a compound of formula (I) according to any one of claims 1
30 to 18 or a pharmaceutically acceptable salt and one or more other therapeutic ingredients.

24. The use of a compound of Formula (I) according to any one of Claims 1 to 18 or a pharmaceutically acceptable salt or physiologically functional derivative thereof in the manufacture of a medicament for the prophylaxis or treatment of a clinical condition for which a selective PDE4 inhibitor is indicated.

5

25. A process for preparing a compound of formula (I) as claimed in claim 1 which process comprises:

(i) treating a compound of formula II:




(II)

10

wherein R^{34} , R^{19} , and R^{20} are as defined in claim 1, and X represents a halogen atom, with an amine of formula R^1R^2NH , wherein R^1 and R^2 are as defined in claim 1;

15

(ii) when $n = 0$, treating a compound of formula (III):



(III)

20


wherein R^1 , R^2 and R^{20} are as defined in claim 1, and Z represents hydrogen, C_{1-6} alkyl or halogen for example chlorine and Y represents hydrogen, chlorine, bromine or iodine, with a thiol of formula R^3SH , or the sodium salt thereof, R^3SNa , wherein R^3 is as defined in claim 1, with the proviso that at least one of Y and Z represent halogen;

(iii) deprotection of protected derivatives of compounds of formula (I);

(iv) when R^{34} represents hydrogen, R^{19} represents $R^3S(=O)_n$, and R^1 , R^2 , R^{20} and n are as defined in claim 1,

hydrogenation of a compound of formula (XXIX)

5



wherein Y represents chlorine, bromine or iodine, $n = 1$ or 2 , and R^1 , R^2 , R^3 and R^{20} are as defined in claim 1.

10

and thereafter and if so desired, converting a first compound of formula (I) into a further compound of formula (I);

and / or thereafter and if so desired converting a compound of formula (I) into a salt thereof.

15

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005494

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/12 C07D215/54 C07D413/12 C07D403/12 C07D407/12
C07D401/12 C07D409/12 C07D403/14 C07D407/14 A61K31/4706
A61K31/4709 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 480 052 A (OTSUKA PHARMA CO LTD) 15 April 1992 (1992-04-15) page 3, line 24 - page 3, line 49; examples 95,96,21	1-15, 20-23,25
X	WO 02/092571 A (ASTRAZENECA AB ; LARSSON JOAKIM (SE); SJOE PETER (SE)) 21 November 2002 (2002-11-21) page 2, line 15 - page 3, line 36; examples 119,120,184e,155,156,184	1-15, 20-23,25
X	WO 02/20489 A (SQUIBB BRISTOL MYERS CO ; YU GUIXUE (US); BI YINGZHI (US); MACOR JOHN) 14 March 2002 (2002-03-14) page 2, line 24 - page 4, line 3	1-15, 20-23,25
A		16-19,24
A	WO 98/57936 A (DARWIN DISCOVERY LTD) 23 December 1998 (1998-12-23) page 2, line 3 - page 3, line 14	1-25

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

25 August 2004

Date of mailing of the international search report

01/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Usueli, A

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2004/005494

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/005494

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0480052	A	15-04-1992	AU 634880 B2	04-03-1993
			AU 7548091 A	21-10-1991
			DE 69128682 D1	19-02-1998
			DE 69128682 T2	04-06-1998
			DK 480052 T3	11-05-1998
			EP 0480052 A1	15-04-1992
			ES 2112272 T3	01-04-1998
			JP 4211661 A	03-08-1992
			WO 9114677 A1	03-10-1991
			US 5215999 A	01-06-1993
WO 02092571	A	21-11-2002	BR 0209431 A	03-08-2004
			CA 2446717 A1	21-11-2002
			CZ 20033034 A3	16-06-2004
			EE 200300544 A	16-02-2004
			EP 1387830 A1	11-02-2004
			WO 02092571 A1	21-11-2002
WO 0220489	A	14-03-2002	AU 8516301 A	22-03-2002
			WO 0220489 A2	14-03-2002
			US 2003225128 A1	04-12-2003
			US 2002177587 A1	28-11-2002
WO 9857936	A	23-12-1998	AU 735573 B2	12-07-2001
			AU 8222898 A	04-01-1999
			CA 2289100 A1	23-12-1998
			CN 1260784 T	19-07-2000
			EP 0993449 A1	19-04-2000
			WO 9857936 A1	23-12-1998
			JP 2002504925 T	12-02-2002
			US 6069151 A	30-05-2000
			ZA 9805253 A	17-06-1999